

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 601 135 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
02.11.2005 Bulletin 2005/44

(51) Int Cl.7: **A61K 38/17, A61K 38/18,
C07K 14/51**

(21) Application number: **93900497.4**

(86) International application number:
PCT/US1992/007432

(22) Date of filing: **28.08.1992**

(87) International publication number:
WO 1993/005751 (01.04.1993 Gazette 1993/09)

(54) OSTEOGENIC PROTEINS IN THE TREATMENT OF METABOLIC BONE DISEASES

OSTEOGENISCHE PROTEINE IN DER BEHANDLUNG VON METABOLISCHEN
KNOCHENKRANKHEITEN

PROTEINES OSTEOGENIQUES POUR LE TRAITEMENT DES MALADIES OSSEUSES
METABOLIQUES

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
SE**

(74) Representative: **Price, Vincent Andrew et al
Fry Heath & Spence LLP
The Gables
Massetts Road
Horley Surrey RH6 7DQ (GB)**

(30) Priority: **30.08.1991 US 752857
30.08.1991 US 752764
30.08.1991 US 752861
31.07.1992 US 923780**

(56) References cited:
**EP-A- 0 416 578 EP-A- 0 512 844
EP-A- 0 514 720 WO-A-89/09788
WO-A-89/10409 WO-A-91/05802
WO-A-92/00382 WO-A-92/15323
WO-A-92/21365**

(43) Date of publication of application:
15.06.1994 Bulletin 1994/24

(73) Proprietor: **Curis, Inc.
Cambridge, MA 02138 (US)**

(72) Inventors:
• **KUBERASAMPATH, Thangavel
Medway, MA 02053 (US)**
• **COHEN, Charles, M.
Medway, MA 02053 (US)**
• **OPPERMANN, Hermann
Medway, MA 02053 (US)**
• **OZKAYNAK, Engin
Milford, MA 01757 (US)**
• **RUEGER, David, C.
Hopkinton, MA 01748 (US)**
• **PANG, Roy, H., L.
Etna, NH 03750 (US)**
• **SMART, John, E.
Weston, MA 02193 (US)**

- **PROCEEDINGS OF THE NATIONAL ACADEMY
OF SCIENCES OF USA. vol. 87, March 1990,
WASHINGTON US pages 2220 - 2224 WANG, E.A.
ET AL 'Recombinant human bone
morphogenetic protein induces bone formation'**
- **PROCEEDINGS OF THE NATIONAL ACADEMY
OF SCIENCES OF USA. vol. 86, November 1989,
WASHINGTON US pages 8793 - 8797
VUKICEVIC, S. ET AL 'Stimulation of the
expression of osteogenic and chondrogenic
phenotypes in vitro by osteogenin'**
- **JOURNAL OF BIOLOGICAL CHEMISTRY.
(MICROFILMS) vol. 267, no. 28, 5 October 1992,
BALTIMORE, MD US pages 20352 - 20362
SAMPATH T.K. ET AL 'Recombinant human
osteogenic protein-1 (hOP-1) induces new bone
formation in vivo with a specific activity
comparable with natural bovine osteogenic
protein and stimulates osteoblast proliferation
and differentiation in vitro'**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 601 135 B1

- EMBO JOURNAL vol. 9, no. 7, 1990, EYNHAM, OXFORD GB pages 2085 - 2093 \ZKAYNAK, E. ET AL 'OP-1 cDNA encodes an osteogenic protein in the TGF-beta family'

Description

[0001] This invention relates to means for increasing the bone mass and/or preventing the loss of bone mass in a mammal. In particular it relates to use of a morphogen in the manufacture of a medicament for treatment of metabolic bone disease.

Background of the Invention

[0002] WO 92/21365 discloses a method for generating new bone growth in a mammal comprising administering to the mammal a safe and effective amount of Vitamin D compound in combination with a safe and effective amount of osteoinductive extract or at least one BMP.

[0003] EP 0 514 720 discloses bone growth factors used to stimulate bone formation when administered with agents that inhibit bone resorption.

[0004] Throughout adult life, bone is continually undergoing remodeling through the interactive cycles of bone formation and resorption (bone turnover). Bone resorption typically is rapid, and is mediated by osteoclasts (bone resorbing cells), formed by mononuclear phagocytic precursor cells at bone remodeling sites. This process then is followed by the appearance of osteoblasts (bone forming cells) which form bone slowly to replace the lost bone. The activities of the various cell types that participate in the remodeling process are controlled by interacting systemic (e.g., hormones, lymphokines, growth factors, vitamins) and local factors (e.g., cytokines, adhesion molecules, lymphokines and growth factors). The fact that completion of this process normally leads to balanced replacement and renewal of bone indicates that the molecular signals and events that influence bone remodeling are tightly controlled.

[0005] A number of bone growth disorders are known which cause an imbalance in the bone remodeling cycle. Chief among these are metabolic bone diseases, such as osteoporosis, osteoplasia (osteomalacia), chronic renal failure and hyperparathyroidism, which result in abnormal or excessive loss of bone mass (osteopenia). Other bone diseases, such as Paget's disease, also cause excessive loss of bone mass at localized sites.

[0006] Osteoporosis is a structural deterioration of the skeleton caused by loss of bone mass resulting from an imbalance in bone formation, bone resorption, or both, such that the resorption dominates the bone formation phase, thereby reducing the weight-bearing capacity of the affected bone. In a healthy adult, the rate at which bone is formed and resorbed is tightly coordinated so as to maintain the renewal of skeletal bone. However, in osteoporotic individuals an imbalance in these bone remodeling cycles develops which results in both loss of bone mass and in formation of microarchitectural defects in the continuity of the skeleton. These skeletal defects, created by perturbation in the remodeling sequence, accumulate and finally reach a point at which the structural integrity of the skeleton is severely compromised and bone fracture is likely. Although this imbalance occurs gradually in most individuals as they age ("senile osteoporosis"), it is much more severe and occurs at a rapid rate in postmenopausal women. In addition, osteoporosis also may result from nutritional and endocrine imbalances, hereditary disorders and a number of malignant transformations.

[0007] Patients suffering from chronic renal (kidney) failure almost universally suffer loss of skeletal bone mass (renal osteodystrophy). While it is known that kidney malfunction causes a calcium and phosphate imbalance in the blood, to date replenishment of calcium and phosphate by dialysis does not significantly inhibit osteodystrophy in patients suffering from chronic renal failure. In adults, osteodystrophic symptoms often are a significant cause of morbidity. In children, renal failure often results in a failure to grow, due to the failure to maintain and/or to increase bone mass.

[0008] Osteoplasia, also known as osteomalacia ("soft bones"), is a defect in bone mineralization (e.g., incomplete mineralization), and classically is related to vitamin D deficiency (1,25-dihydroxy vitamin D₃). The defect can cause compression fractures in bone, and a decrease in bone mass, as well as extended zones of hypertrophy and proliferative cartilage in place of bone tissue. The deficiency may result from a nutritional deficiency (e.g., rickets in children), malabsorption of vitamin D or calcium, and/or impaired metabolism of the vitamin.

[0009] Hyperparathyroidism (overproduction of the parathyroid hormone) is known to cause malabsorption of calcium, leading to abnormal bone loss. In children, hyperparathyroidism can inhibit growth, in adults the skeleton integrity is compromised and fracture of the ribs and vertebrae are characteristic. The parathyroid hormone imbalance typically may result from thyroid adenomas or gland hyperplasia, or may result from prolonged pharmacological use of a steroid. Secondary hyperparathyroidism also may result from renal osteodystrophy. In the early stages of the disease osteoclasts are stimulated to resorb bone in response to the excess hormone present. As the disease progresses, the trabecular bone ultimately is resorbed and marrow is replaced with fibrosis, macrophages and areas of hemorrhage as a consequence of microfractures. This condition is referred to clinically as osteitis fibrosa.

[0010] Paget's disease (osteitis deformans) is a disorder currently thought to have a viral etiology and is characterized by excessive bone resorption at localized sites which flare and heal but which ultimately are chronic and progressive, and may lead to malignant transformation. The disease typically affects adults over the age of 25.

[0011] To date, osteopenia treatments are based on inhibiting further bone resorption, e.g., by 1) inhibiting the dif-

ferentiation of hemopoietic mononuclear cells into mature osteoclasts, 2) by directly preventing osteoclast-mediated bone resorption, or 3) by affecting the hormonal control of bone resorption. Drug regimens used for the treatment of osteoporosis include calcium supplements, estrogen, calcitonin and diphosphonates. Vitamin D₃ and its metabolites, known to enhance calcium and phosphate absorption, also are being tried. None of the current therapies stimulate regeneration of new bone tissue. In addition, all of these agents have only a transient effect on bone remodeling. Thus, while in some cases the progression of the disease may be halted or slowed, patients with significant bone deterioration remain actively at risk. This is particularly prevalent in disorders such as osteoporosis where early diagnosis is difficult and/or rare and significant structural deterioration of the bone already may have occurred.

[0012] It is an object of the present invention to develop medical use of a morphogen for treatment of metabolic bone disease in an individual who, for example, is afflicted with a disease which decreases skeletal bone mass, particularly where the disease causes an imbalance in bone remodeling. Another object is use to enhance bone growth in children suffering from bone disorders, including metabolic bone diseases. Still another object is to prevent or inhibit bone deterioration in individuals at risk for loss of bone mass, including postmenopausal women, aged individuals, and patients undergoing dialysis. Yet another object is to provide methods and compositions for repairing defects in the microstructure of structurally compromised bone, including repairing bone fractures. Thus, the invention is aimed at stimulating bone formation and increasing bone mass, optionally over prolonged periods of time, and particularly to decrease the occurrence of new fractures resulting from structural deterioration of the skeleton. These and other objects and features of the invention will be apparent from the description, drawings, and claims which follow.

Summary of the Invention

[0013] The present invention provides use of a morphogen as defined in the appended claims.

[0014] In one aspect, the invention features use of a morphogen in the manufacture of a medicament for treatment of metabolic bone disease in a mammal. The treatment includes administering to the individual a therapeutically effective morphogen in an amount and for a time sufficient to inhibit the loss of bone mass, and/or to increase bone mass in the individual.

[0015] A therapeutic treatment method and composition for preventing loss of bone mass and/or for increasing bone mass in a mammal could include administering to the mammal a compound that stimulates *in vivo* a therapeutically effective concentration of an endogenous morphogen in the body of the mammal sufficient to prevent loss of and/or to increase bone mass in the individual. These compounds are referred to herein as morphogen-stimulating agents, and are understood to include substances which, when administered to a mammal, act on tissue(s) or organ(s) that normally are responsible for, or capable of, producing a morphogen and/or secreting a morphogen, and which cause the endogenous level of the morphogen to be altered. The agent may act, for example, by stimulating expression and/or secretion of an endogenous morphogen.

[0016] The morphogens described herein are believed to play a significant role in maintaining appropriate bone mass in an individual. Thus, a morphogen may be administered to any individual who requires assistance in maintaining appropriate bone mass and/or who suffers from a bone remodeling imbalance. For example, the morphogen or morphogen-stimulating agent may be administered to an adult suffering from renal failure to prevent bone deterioration which is associated with that disease, e.g., to correct bone loss due to late stage kidney failure. Similarly, the administration of a morphogen to a child suffering from renal failure is expected not only to alleviate loss of bone mass in the child, but also to stimulate bone formation and thus growth. In addition, administration of a morphogen or morphogen-stimulating agent to an individual suffering from defects in skeletal microstructure is expected to result in repair of that defect, and to enhance the weight-bearing capacity of the treated bone.

[0017] Accordingly, treatment methods and compositions may be used to treat a bone fracture or any disease which causes or results in bone fractures or other defects in skeletal microstructure, including loss of bone mass, and which compromise the weight-bearing capacity of bone. Such diseases include, for example, chronic renal failure and other kidney diseases, particularly those requiring dialysis; osteomalacia; vitamin D deficiency-induced osteopenia or osteoporosis; postmenopausal or senile osteoporosis; hyperparathyroidism and Paget's disease.

[0018] A morphogen or morphogen-stimulating agent could be administered systemically to the individual, e.g., orally or parenterally. In another embodiment the morphogen may be provided directly to the bone, e.g., by injection to the bone periosteum or endosteum. Direct injection is particularly useful for repairing defects in the microstructure of the bone, including bone fractures.

[0019] In any treatment method "administration of morphogen" refers to the administration of the morphogen, either alone or in combination with other molecules. For example, the mature form of the morphogen may be provided in association with its precursor "pro" domain, which is known to enhance the solubility of the protein. Other useful molecules known to enhance protein solubility include casein and other milk components, as well as various serum proteins. Additional useful molecules which may be associated with the morphogen or morphogen-stimulating agent include tissue targeting molecules capable of directing the morphogen or morphogen-stimulating agent to bone. Tissue tar-

getting molecules envisioned to be useful in the treatment protocols include tetracycline, diphosphonates, and antibodies or other binding proteins which interact specifically with surface molecules on bone tissue cells.

[0020] Still another useful tissue targeting molecule is the morphogen precursor "pro" domain, particularly that of OP-1, BMP2 or BMP4. These proteins are found naturally associated with bone tissue but likely are synthesized in other tissues and targeted to bone tissue after secretion from the synthesizing tissue. For example, the primary source of OP-1 synthesis appears to be the tissue of the urinary tract (e.g., renal tissue), while the protein has been shown to be active in bone tissue (see below.) Moreover, the protein has been identified in serum, saliva and various milk forms. In addition, the secreted form of the protein comprises the mature dimer in association with the pro domain of the intact morphogen sequence. Accordingly, the associated morphogen pro domains may act to target specific morphogens to different tissues in vivo.

[0021] Associated tissue targeting or solubility-enhancing molecules also may be covalently linked to the morphogen using standard chemical means, including acid-labile linkages, which likely will be preferentially cleaved in the acidic environment of bone remodeling sites.

[0022] The morphogens or morphogen-stimulating agents also may be administered together with other "co-factors" known to have a beneficial effect on bone remodeling, including parathyroid hormone, vitamin D₃, prostaglandins, dexamethasone, IGF (I, II) and their binding proteins, and other agents known to enhance osteoblast activity. Other useful cofactors include calcitonin and estrogen and other agents which inhibit bone resorption.

[0023] Among the morphogens useful in this invention are proteins originally identified as osteogenic proteins, such as the OP-1, OP-2 and CHMP2 proteins, as well as amino acid sequence-related proteins such as DPP (from *Drosophila*), Vgl (from *Xenopus*), Vgr-1 (from mouse, see U.S. 5,011,691 to Oppermann et al.), GDF-1 (from mouse, see Lee (1991) *PNAS* 88:4250-4254), all of which are presented in Table II and Seq. ID Nos. 5-14), and the recently identified 60A protein (from *Drosophila*, Seq. ID No. 24, see Wharton et al. (1991) *PNAS* 88:9214-9218.) The members of this family, which include members of the TGF- β super-family of proteins, share substantial amino acid sequence homology in their C-terminal regions. The proteins are translated as a precursor, having an N-terminal signal peptide sequence, typically less than about 30 residues, followed by a "pro" domain that is cleaved to yield the mature sequence. The signal peptide is cleaved rapidly upon translation, at a cleavage site that can be predicted in a given sequence using the method of Von Heijne ((1986) *Nucleic Acids Research* 14:4683-4691.) Table I, below, describes the various morphogens identified to date, including their nomenclature as used herein, their Seq. ID references, and publication sources for the amino acid sequences of the full length proteins not included in the Seq. Listing.

TABLE I

"OP-1"	Refers generically to the group of morphogenically active proteins expressed from part or all of a DNA sequence encoding OP-1 protein, including allelic and species variants thereof, e.g., human OP-1 ("hOP-1", Seq. ID No. 5, mature protein amino acid sequence), or mouse OP-1 ("mOP-1", Seq. ID No. 6, mature protein amino acid sequence.) The conserved seven cysteine skeleton is defined by residues 38 to 139 of Seq. ID Nos. 5 and 6. The cDNA sequences and the amino acids encoding the full length proteins are provided in Seq. ID Nos. 16 and 17 (hOP1) and Seq. ID Nos. 18 and 19 (mOP1.) The mature proteins are defined by residues 293-431 (hOP1) and 292-430 (mOP1). The "pro" regions of the proteins, cleaved to yield the mature, morphogenically active proteins are defined essentially by residues 30-292 (hOP1) and residues 30-291 (mOP1).
"OP-2"	refers generically to the group of active proteins expressed from part or all of a DNA sequence encoding OP-2 protein, including allelic and species variants thereof, e.g., human OP-2 ("hOP-2", Seq. ID No. 7, mature protein amino acid sequence) or mouse OP-2 ("mOP-2", Seq. ID No. 8, mature protein amino acid sequence). The conserved seven cysteine skeleton is defined by residues 38 to 139 of Seq. ID Nos. 7 and 8. The cDNA sequences and the amino acids encoding the full length proteins are provided in Seq. ID Nos. 20 and 21 (hOP2) and Seq. ID Nos. 22 and 23 (mOP2.) The mature proteins are defined essentially by residues 264-402 (hOP2) and 261-399 (mOP2). The "pro" regions of the proteins, cleaved to yield the mature, morphogenically active proteins are defined essentially by residues 18-263 (hOP2) and residues 18-260 (mOP2). (Another cleavage site also occurs 21 residues further upstream for both OP-2 proteins.)

TABLE I (continued)

5	"CBMP2"	refers generically to the morphogenically active proteins expressed from a DNA sequence encoding the CBMP2 proteins, including allelic and species variants thereof, e.g., human CBMP2A ("CBMP2A (fx)", Seq ID No. 9) or human CBMP2B DNA ("CBMP2B(fx)", Seq. ID No. 10). The amino acid sequence for the full length proteins, referred to in the literature collectively as BMP2A and BMP2B, or BMP2 and BMP4, appear in Wozney, et al. (1988) <u>Science</u> 242:1528-1534. The pro domain for BMP2 (BMP2A) likely includes residues 25-248 or 25-282; the mature protein, residues 249-396 or 283-396. The pro domain for BMP4 (BMP2B) likely includes residues 25-256 or 25-292; the mature protein, residues 257-408 or 293-408.
10	"DPP(fx)"	refers to protein sequences encoded by the Drosophila DPP gene and defining the conserved seven cysteine skeleton (Seq. ID No. 11). The amino acid sequence for the full length protein appears in Padgett, et al (1987) <u>Nature</u> 325: 81-84. The pro domain likely extends from the signal peptide cleavage site to residue 456; the mature protein likely is defined by residues 457-588.
15	"Vgl(fx)"	refers to protein sequences encoded by the Xenopus Vgl gene and defining the conserved seven cysteine skeleton (Seq. ID No. 12). The amino acid sequence for the full length protein appears in Weeks (1987) <u>Cell</u> 51: 861-867. The pro domain likely extends from the signal peptide cleavage site to residue 246; the mature protein likely is defined by residues 247-360.
20	"Vgr-1(fx)"	refers to protein sequences encoded by the murine Vgr-1 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 13). The amino acid sequence for the full length protein appears in Lyons, et al, (1989) <u>PNAS</u> 86: 4554-4558. The pro domain likely extends from the signal peptide cleavage site to residue 299; the mature protein likely is defined by residues 300-438.
25	"GDF-1(fx)"	refers to protein sequences encoded by the human GDF-1 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 14). The cDNA and encoded amino sequence for the full length protein is provided in Seq. ID. No. 32. The pro domain likely extends from the signal peptide cleavage site to residue 214; the mature protein likely is defined by residues 215-372.
30	"60A"	refers generically to the morphogenically active proteins expressed from part or all of a DNA sequence (from the Drosophila 60A gene) encoding the 60A proteins (see Seq. ID No. 24 wherein the cDNA and encoded amino acid sequence for the full length protein is provided). "60A(fx)" refers to the protein sequences defining the conserved seven cysteine skeleton (residues 354 to 455 of Seq. ID No. 24.) The pro domain likely extends from the signal peptide cleavage site to residue 324; the mature protein likely is defined by residues 325-455.
35	"BMP3(fx)"	refers to protein sequences encoded by the human BMP3 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 26). The amino acid sequence for the full length protein appears in wozney et al. (1988) <u>Science</u> 242: 1528-1534. The pro domain likely extends from the signal peptide cleavage site to residue 290; the mature protein likely is defined by residues 291-472.
40	"BMP5(fx)"	refers to protein sequences encoded by the human BMP5 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 27). The amino acid sequence for the full length protein appears in Celeste, et al. (1991) <u>PNAS</u> 87: 9843-9847. The pro domain likely extends from the signal peptide cleavage site to residue 316; the mature protein likely is defined by residues 317-454.
45	"BMP6(fx)"	refers to protein sequences encoded by the human BMP6 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 28). The amino acid sequence for the full length protein appear sin Celeste, et al. (1990) <u>PNAS</u> 87: 9843-5847. The pro domain likely includes extends from the signal peptide cleavage site to residue 374; the mature sequence likely includes residues 375-513.

[0024] The OP-2 proteins have an additional cysteine residue in this region (e.g., see residue 41 of Seq. ID Nos. 7 and 8), in addition to the conserved cysteine skeleton in common with the other proteins in this family. The GDF-1 protein has a four amino acid insert within the conserved skeleton (residues 44-47 of Seq. ID No. 14) but this insert likely does not interfere with the relationship of the cysteines in the folded structure. In addition, the CBMP2 proteins are missing one amino acid residue within the cysteine skeleton.

[0025] The morphogens are inactive when reduced, but are active as oxidized homodimers and when oxidized in combination with other morphogens of this invention. Thus, as defined herein, a morphogen is a dimeric protein comprising a pair of polypeptide chains, wherein each polypeptide chain comprises at least the C-terminal six cysteine skeleton defined by residues 43-139 of Seq. ID No. 5, including functionally equivalent arrangements of these cysteines (e.g., amino acid insertions or deletions which alter the linear arrangement of the cysteines in the sequence but not their relationship in the folded structure), such that, when the polypeptide chains are folded, the dimeric protein species comprising the pair of polypeptide chains has the appropriate three-dimensional structure, including the appropriate

intra- or inter-chain disulfide bonds such that the protein is capable of acting as a morphogen as defined herein. Specifically, the morphogens generally are capable of all of the following biological functions in a morphogenically permissive environment: stimulating proliferation of progenitor cells; stimulating the differentiation of progenitor cells; stimulating the proliferation of differentiated cells; and supporting the growth and maintenance of differentiated cells, including the "redifferentiation" of transformed cells. In addition, it also is anticipated that these morphogens are capable of inducing redifferentiation of committed cells under appropriate environmental conditions.

[0026] The morphogens used in this invention comprise one of two species of generic amino acid sequences: Generic Sequence 1 (Seq. ID No. 1) or Generic Sequence 2 (Seq. ID No. 2); where each Xaa indicates one of the 20 naturally-occurring L-isomer, α -amino acids or a derivative thereof. Generic Sequence 1 comprises the conserved six cysteine skeleton and Generic Sequence 2 comprises the conserved six cysteine skeleton plus the additional cysteine identified in OP-2 (see residue 36, Seq. ID No. 2). In another preferred aspect, these sequences further comprise the following additional sequence at their N-terminus:

Cys Xaa Xaa Xaa Xaa (Seq. ID No. 15)
1 5

[0027] Preferred amino acid sequences within the foregoing generic sequences include: Generic Sequence 3 (Seq. ID No. 3), Generic Sequence 4 (Seq. ID No. 4), Generic Sequence 5 (Seq. ID No. 30) and Generic Sequence 6 (Seq. ID No. 31), listed below. These Generic Sequences accommodate the homologies shared among the various preferred members of this morphogen family identified in Table II, as well as the amino acid sequence variation among them. Specifically, Generic Sequences 3 and 4 are composite amino acid sequences of the following proteins presented in Table II and identified in Seq. ID Nos. 5-14: human OP-1 (hOP-1, Seq. ID Nos. 5 and 16-17), mouse OP-1 (mOP-1, Seq. ID Nos. 6 and 18-19), human and mouse OP-2 (Seq. ID Nos. 7, 8, and 20-22), CBMP2A (Seq. ID No. 9), CBMP2B (Seq. ID No. 10), DPP (from *Drosophila*, Seq. ID No. 11), Vgl, (from *Xenopus*, Seq. ID No. 12), Vgr-1 (from mouse, Seq. ID No. 13), and GDF-1 (from mouse, Seq. ID No. 14.) The generic sequences include both the amino acid identity shared by the sequences in Table II, as well as alternative residues for the variable positions within the sequence. Note that these generic sequences allow for an additional cysteine at position 41 or 46 in Generic Sequences 3 or 4, respectively, providing an appropriate cysteine skeleton where inter- or intramolecular disulfide bonds can form, and contain certain critical amino acids which influence the tertiary structure of the proteins.

Generic Sequence 3

[0028]

5 Leu Tyr Val Xaa Phe
 1 5
 10 Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
 10
 Xaa Ala Pro Xaa Gly Xaa Xaa Ala
 15 15 20
 Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
 25 30
 20 Xaa Pro Xaa Xaa Xaa Xaa Xaa
 35
 25 Xaa Xaa Xaa Asn His Ala Xaa Xaa
 40 45
 Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa
 30 50
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
 55 60
 35 Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
 65
 40
 Xaa Xaa Xaa Leu Xaa Xaa Xaa
 45 70 75
 Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
 80
 50 Xaa Xaa Xaa Xaa Met Xaa Val Xaa
 85 90
 55 Xaa Cys Gly Cys Xaa
 95

wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows: "Res." means "residue" and Xaa at res.4 = (Ser, Asp or Glu); Xaa at res.6 = (Arg, Gln, Ser or Lys); Xaa at res.7 = (Asp or Glu); Xaa at res.8 = (Leu or Val); Xaa at res.11 = (Gln, Leu, Asp, His or Asn); Xaa at res.12 = (Asp, Arg or Asn); Xaa at res.14 = (Ile or Val); Xaa at res.15 = (Ile or Val); Xaa at res.18 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.20 = (Tyr or Phe); Xaa at res.21 = (Ala, Ser, Asp, Met, His, Leu or Gln); Xaa at res.23 = (Tyr, Asn or Phe); Xaa at res.26 = (Glu, His, Tyr, Asp or Gln); Xaa at res.28 = (Glu, Lys, Asp or Gln); Xaa at res.30 = (Ala, Ser, Pro or Gln); Xaa at res.31 = (Phe, Leu or Tyr); Xaa at res.33 = (Leu or Val); Xaa at res.34 = (Asn, Asp, Ala or Thr); Xaa at res.35 = (Ser, Asp, Glu, Leu or Ala); Xaa at res.36 = (Tyr, Cys, His, Ser or Ile); Xaa at res.37 = (Met, Phe, Gly or Leu); Xaa at res.38 = (Asn or Ser); Xaa at res.39 = (Ala, Ser or Gly); Xaa at res.40 = (Thr, Leu or Ser); Xaa at res.44 = (Ile or Val); Xaa at res.45 = (Val or Leu); Xaa at res.46 = (Gln or Arg); Xaa at res.47 = (Thr, Ala or Ser); Xaa at res.49 = (Val or Met); Xaa at res.50 = (His or Asn); Xaa at res.51 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.52 = (Ile, Met, Asn, Ala or Val); Xaa at res.53 = (Asn, Lys, Ala or Glu); Xaa at res.54 = (Pro or Ser); Xaa at res.55 = (Glu, Asp, Asn, or Gly); Xaa at res.56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser or Ala); Xaa at res.57 = (Val, Ala or Ile); Xaa at res.58 = (Pro or Asp); Xaa at res.59 = (Lys or Leu); Xaa at res.60 = (Pro or Ala); Xaa at res.63 = (Ala or Val); Xaa at res.65 = (Thr or Ala); Xaa at res.66 = (Gln, Lys, Arg or Glu); Xaa at res.67 = (Leu, Met or Val); Xaa at res.68 = (Asn, Ser or Asp); Xaa at res.69 = (Ala, Pro or Ser); Xaa at res.70 = (Ile, Thr or Val); Xaa at res.71 = (Ser or Ala); Xaa at res.72 = (Val or Met); Xaa at res.74 = (Tyr or Phe); Xaa at res.75 = (Phe, Tyr or Leu); Xaa at res.76 = (Asp or Asn); Xaa at res.77 = (Asp, Glu, Asn or Ser); Xaa at res.78 = (Ser, Gln, Asn or Tyr); Xaa at res.79 = (Ser, Asn, Asp or Glu); Xaa at res.80 = (Asn, Thr or Lys); Xaa at res.82 = (Ile or Val); Xaa at res.84 = (Lys or Arg); Xaa at res.85 = (Lys, Asn, Gln or His); Xaa at res.86 = (Tyr or His); Xaa at res.87 = (Arg, Gln or Glu); Xaa at res.88 = (Asn, Glu or Asp); Xaa at res.90 = (Val, Thr or Ala); Xaa at res.92 = (Arg, Lys, Val, Asp or Glu); Xaa at res.93 = (Ala, Gly or Glu); and Xaa at res.97 = (His or Arg);

Generic Sequence 4

[0029]

```

Cys Xaa Xaa Xaa Xaa Leu Tyr Val Xaa Phe
  1                      5                      10
Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
                15
Xaa Ala Pro Xaa Gly Xaa Xaa Ala
  20                      25
Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
                30                      35

```

Xaa Pro Xaa Xaa Xaa Xaa Xaa
 40
 Xaa Xaa Xaa Asn His Ala Xaa Xaa
 45 50
 Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa
 55
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
 60 65
 Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
 70
 Xaa Xaa Xaa Leu Xaa Xaa Xaa
 75 80
 Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
 85
 Xaa Xaa Xaa Xaa Met Xaa Val Xaa
 90 95
 Xaa Cys Gly Cys Xaa
 100

wherein each Xaa is independently selected from a group of one or more specified amino acids as defined by the following: "Res." means "residue" and Xaa at res.2 = (Lys or Arg); Xaa at res.3 = (Lys or Arg); Xaa at res.4 = (His or Arg); Xaa at res.5 = (Glu, Ser, His, Gly, Arg or Pro); Xaa at res.9 = (Ser, Asp or Glu); Xaa at res.11 = (Arg, Gln, Ser or Lys); Xaa at res.12 = (Asp or Glu); Xaa at res.13 = (Leu or Val); Xaa at res.16 = (Gln, Leu, Asp, His or Asn); Xaa at res.17 = (Asp, Arg, or Asn); Xaa at res.19 = (Ile or Val); Xaa at res.20 = (Ile or Val); Xaa at res.23 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.25 = (Tyr or Phe); Xaa at res.26 = (Ala, Ser, Asp, Met, His, Leu, or Gln); Xaa at res.28 = (Tyr, Asn or Phe); Xaa at res.31 = (Glu, His, Tyr, Asp or Gln); Xaa at res.33 = Glu, Lys, Asp or Gln; Xaa at res.35 = (Ala, Ser or Pro); Xaa at res.36 = (Phe, Leu or Tyr); Xaa at res.38 = (Leu or Val); Xaa at res.39 = (Asn, Asp, Ala or Thr); Xaa at res.40 = (Ser, Asp, Glu, Leu or Ala); Xaa at res.41 = (Tyr, Cys, His, Ser or Ile); Xaa at res.42 = (Met, Phe, Gly or Leu); Xaa at res.44 = (Ala, Ser or Gly); Xaa at res.45 = (Thr, Leu or Ser); Xaa at res.49 = (Ile or Val); Xaa at res.50 = (Val or Leu); Xaa at res.51 = (Gln or Arg); Xaa at res.52 = (Thr, Ala or Ser); Xaa at res.54 = (Val or Met); Xaa at res.55 = (His or Asn); Xaa at res.56 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.57 = (Ile, Met, Asn, Ala or Val); Xaa at res.58 = (Asn, Lys, Ala or Glu); Xaa at res.59 = (Pro or Ser); Xaa at res.60 = (Glu, Asp, or Gly); Xaa at res.61 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser or Ala); Xaa at res.62 = (Val, Ala or Ile); Xaa at res.63 = (Pro or Asp); Xaa at res.64 = (Lys or Leu); Xaa at res.65 = (Pro or Ala); Xaa at res.68 = (Ala or Val); Xaa at res.70 = (Thr or Ala); Xaa at res.71 = (Gln, Lys, Arg or Glu); Xaa at res.72 = (Leu, Met or Val); Xaa at res.73 = (Asn, Ser or Asp); Xaa at res.74 = (Ala, Pro or Ser); Xaa at res.75 = (Ile, Thr or Val); Xaa at res.76 = (Ser or Ala); Xaa at res.77 = (Val or Met); Xaa at res.79 = (Tyr or Phe); Xaa at res.80 = (Phe, Tyr or Leu); Xaa at res.81 = (Asp or Asn); Xaa at res.82 = (Asp, Glu, Asn or Ser); Xaa at res.83 = (Ser, Gln, Asn or Tyr); Xaa at res.84 = (Ser, Asn, Asp or Glu); Xaa at res.85 = (Asn, Thr or Lys); Xaa at res.87 = (Ile or Val); Xaa at res.89 = (Lys or Arg); Xaa at res.90 = (Lys, Asn, Gln or His); Xaa at res.91 = (Tyr or His); Xaa at res.92 = (Arg, Gln or Glu); Xaa at res.93 = (Asn, Glu or Asp); Xaa at res.95 = (Val, Thr or Ala); Xaa at res.97 = (Arg, Lys, Val, Asp or Glu); Xaa at res.98 = (Ala, Gly or Glu); and Xaa at res.102 = (His or Arg).

[0030] Similarly, Generic Sequence 5 (Seq. ID No. 30) and Generic Sequence 6 (Seq. ID No. 31) accommodate the homologies shared among all the morphogen protein family members identified in Table II. Specifically, Generic Sequences 5 and 6 are composite amino acid sequences of human OP-1 (hOP-1, Seq. ID Nos. 5 and 16-17), mouse OP-1 (mOP-1, Seq. ID Nos. 6 and 18-19), human and mouse OP-2 (Seq. ID Nos. 7, 8, and 20-22), CBMP2A (Seq. ID No. 9), CBMP2B (Seq. ID No. 10), DPP (from Drosophila, Seq. ID No. 11), Vgl, (from Xenopus, Seq. ID No. 12), Vgr-1 (from mouse, Seq. ID No. 13), and GDF-1 (from mouse, Seq. ID No. 14 and 32), human BMP3 (Seq. ID No. 26), human BMP5 (Seq. ID No. 27), human BMP6 (Seq. ID No. 28) and 60A (from Drosophila, Seq. ID No. 24). The generic sequences include both the amino acid identity shared by these sequences in the C-terminal domain, defined by the six and seven cysteine skeletons (Generic Sequences 5 and 6, respectively), as well as alternative residues for the

variable positions within the sequence. As for Generic Sequences 3 and 4, Generic Sequences 5 and 6 allow for an additional cysteine at position 41 (Generic Sequence 5) or position 46 (Generic Sequence 6), providing an appropriate cysteine skeleton where inter- or intramolecular disulfide bonds can form, and containing certain critical amino acids which influence the tertiary structure of the proteins.

5

Generic Sequence 5**[0031]**

10

Leu Xaa Xaa Xaa Phe

1

5

15

Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa

10

20

25

30

35

40

45

50

55

5 Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala
 15 20
 Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
 25 30
 10 Xaa Pro Xaa Xaa Xaa Xaa Xaa
 35
 Xaa Xaa Xaa Asn His Ala Xaa Xaa
 15 40 45
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 20 50
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
 55 60
 25 Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
 65
 Xaa Xaa Xaa Leu Xaa Xaa Xaa
 30 70 75
 Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
 35 80
 Xaa Xaa Xaa Xaa Met Xaa Val Xaa
 85 90
 40 Xaa Cys Xaa Cys Xaa
 95

45 wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows:
 "Res." means "residue" and Xaa at res.2 = (Tyr or Lys); Xaa at res.3 = Val or Ile); Xaa at res.4 = (Ser, Asp or Glu); Xaa
 at res.6 = (Arg, Gln, Ser, Lys or Ala); Xaa at res.7 = (Asp, Glu or Lys); Xaa at res.8 = (Leu, Val or Ile); Xaa at res.11 =
 (Gln, Leu, Asp, His, Asn or Ser); Xaa at res.12 = (Asp, Arg, Asn or Glu); Xaa at res.14 = (Ile or Val); Xaa at res.15 =
 (Ile or Val); Xaa at res.16 (Ala or Ser); Xaa at res.18 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.19 = (Gly or Ser);
 50 Xaa at res.20 = (Tyr or Phe); Xaa at res.21 = (Ala, Ser, Asp, Met, His, Gln, Leu or Gly); Xaa at res.23 = (Tyr, Asn or
 Phe); Xaa at res.26 = (Glu, His, Tyr, Asp, Gln or Ser); Xaa at res.28 = (Glu, Lys, Asp, Gln or Ala); Xaa at res.30 = (Ala,
 Ser, Pro, Gln or Asn); Xaa at res.31 = (Phe, Leu or Tyr); Xaa at res.33 = (Leu, Val or Met); Xaa at res.34 = (Asn, Asp,
 Ala, Thr or Pro); Xaa at res.35 = (Ser, Asp, Glu, Leu, Ala or Lys); Xaa at res.36 = (Tyr, Cys, His, Ser or Ile); Xaa at res.
 37 = (Met, Phe, Gly or Leu); Xaa at res.38 = (Asn, Ser or Lys); Xaa at res.39 = (Ala, Ser, Gly or Pro); Xaa at res.40 =
 55 (Thr, Leu or Ser); Xaa at res.44 = (Ile, Val or Thr); Xaa at res.45 = (Val, Leu or Ile); Xaa at res.46 = (Gln or Arg); Xaa
 at res.47 = (Thr, Ala or Ser); Xaa at res.48 = (Leu or Ile); Xaa at res.49 = (Val or Met); Xaa at res.50 = (His, Asn or
 Arg); Xaa at res.51 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.52 = (Ile, Met, Asn, Ala, Val or Leu); Xaa at res.53 =
 (Asn, Lys, Ala, Glu, Gly or Phe); Xaa at res.54 = (Pro, Ser or Val); Xaa at res.55 = (Glu, Asp, Asn, Gly, Val or Lys); Xaa

at res.56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Ala, Pro or His); Xaa at res.57 = (Val, Ala or Ile); Xaa at res.58 = (Pro or Asp); Xaa at res.59 = (Lys, Leu or Glu); Xaa at res.60 = (Pro or Ala); Xaa at res.63 = (Ala or Val); Xaa at res.65 = (Thr, Ala or Glu); Xaa at res.66 = (Gln, Lys, Arg or Glu); Xaa at res.67 = (Leu, Met or Val); Xaa at res.68 = (Asn, Ser, Asp or Gly); Xaa at res.69 = (Ala, Pro or Ser); Xaa at res.70 = (Ile, Thr, Val or Leu); Xaa at res.71 = (Ser, Ala or Pro); Xaa at res.72 = (Val, Met or Ile); Xaa at res.74 = (Tyr or Phe); Xaa at res.75 = (Phe, Tyr, Leu or His); Xaa at res.76 = (Asp, Asn or Leu); Xaa at res.77 = (Asp, Glu, Asn or Ser); Xaa at res.78 = (Ser, Gln, Asn, Tyr or Asp); Xaa at res.79 = (Ser, Asn, Asp, Glu or Lys); Xaa at res.80 = (Asn, Thr or Lys); Xaa at res.82 = (Ile, Val or Asn); Xaa at res.84 = (Lys or Arg); Xaa at res.85 = (Lys, Asn, Gln, His or Val); Xaa at res.86 = (Tyr or His); Xaa at res.87 = (Arg, Gln, Glu or Pro); Xaa at res.88 = (Asn, Glu or Asp); Xaa at res.90 = (Val, Thr, Ala or Ile); Xaa at res.92 = (Arg, Lys, Val, Asp or Glu); Xaa at res.93 = (Ala, Gly, Glu or Ser); Xaa at res.95 = (Gly or Ala) and Xaa at res.97 = (His or Arg).

Generic Sequence 6

[0032]

15

Cys Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Phe
1 5 10

20

Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
15

25

Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala
20 25

Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
30 35

30

Xaa Pro Xaa Xaa Xaa Xaa Xaa
40

Xaa Xaa Xaa Asn His Ala Xaa Xaa
45 50

35

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
55

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
60 65

40

Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
70

45

Xaa Xaa Xaa Leu Xaa Xaa Xaa
75 80

Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
85

50

Xaa Xaa Xaa Xaa Met Xaa Val Xaa
90 95

Xaa Cys Xaa Cys Xaa
100

55

wherein each Xaa is independently selected from a group of one or more specified amino acids as defined by the following: "Res." means "residue" and Xaa at res.2 = (Lys, Arg, Ala or Gln); Xaa at res.3 = (Lys, Arg or Met); Xaa at

res.4 = (His, Arg or Gln); Xaa at res.5 = (Glu, Ser, His, Gly, Arg, Pro, Thr, or Tyr); Xaa at res.7 = (Tyr or Lys); Xaa at res.8 = (Val or Ile); Xaa at res.9 = (Ser, Asp or Glu); Xaa at res.11 = (Arg, Gln, Ser, Lys or Ala); Xaa at res.12 = (Asp, Glu, or Lys); Xaa at res.13 = (Leu, Val or Ile); Xaa at res.16 = (Gln, Leu, Asp, His, Asn or Ser); Xaa at res.17 = (Asp, Arg, Asn or Glu); Xaa at res.19 = (Ile or Val); Xaa at res.20 = (Ile or Val); Xaa at res.21 = (Ala or Ser); Xaa at res.23 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.24 = (Gly or Ser); Xaa at res.25 = (Tyr or Phe); Xaa at res.26 = (Ala, Ser, Asp, Met, His, Gln, Leu, or Gly); Xaa at res.28 = (Tyr, Asn or Phe); Xaa at res.31 = (Glu, His, Tyr, Asp, Gln or Ser); Xaa at res.33 = Glu, Lys, Asp, Gln or Ala; Xaa at res.35 = (Ala, Ser, Pro, Gln or Asn); Xaa at res.36 = (Phe, Leu or Tyr); Xaa at res.38 = (Leu, Val or Met); Xaa at res.39 = (Asn, Asp, Ala, Thr or Pro); Xaa at res.40 = (Ser, Asp, Glu, Leu, Ala or Lys); Xaa at res.41 = (Tyr, Cys, His, Ser or Ile); Xaa at res.42 = (Met, Phe, Gly or Leu); Xaa at res.43 = (Asn, Ser or Lys); Xaa at res.44 = (Ala, Ser, Gly or Pro); Xaa at res.45 = (Thr, Leu or Ser); Xaa at res.49 = (Ile, Val or Thr); Xaa at res.50 = (Val, Leu or Ile); Xaa at res.51 = (Gln or Arg); Xaa at res.52 = (Thr, Ala or Ser); Xaa at res.53 = (Leu or Ile); Xaa at res.54 = (Val or Met); Xaa at res.55 = (His, Asn or Arg); Xaa at res.56 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.57 = (Ile, Met, Asn, Ala, Val or Leu); Xaa at res.58 = (Asn, Lys, Ala, Glu, Gly or Phe); Xaa at res.59 = (Pro, Ser or Val); Xaa at res.60 = (Glu, Asp, Gly, Val or Lys); Xaa at res.61 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Ala, Pro or His); Xaa at res.62 = (Val, Ala or Ile); Xaa at res.63 = (Pro or Asp); Xaa at res.64 = (Lys, Leu or Glu); Xaa at res.65 = (Pro or Ala); Xaa at res.68 = (Ala or Val); Xaa at res.70 = (Thr, Ala or Glu); Xaa at res.71 = (Gln, Lys, Arg or Glu); Xaa at res.72 = (Leu, Met or Val); Xaa at res.73 = (Asn, Ser, Asp or Gly); Xaa at res.74 = (Ala, Pro or Ser); Xaa at res.75 = (Ile, Thr, Val or Leu); Xaa at res.76 = (Ser, Ala or Pro); Xaa at res.77 = (Val, Met or Ile); Xaa at res.79 = (Tyr or Phe); Xaa at res.80 = (Phe, Tyr, Leu or His); Xaa at res.81 = (Asp, Asn or Leu); Xaa at res.82 = (Asp, Glu, Asn or Ser); Xaa at res.83 = (Ser, Gln, Asn, Tyr or Asp); Xaa at res.84 = (Ser, Asn, Asp, Glu or Lys); Xaa at res.85 = (Asn, Thr or Lys); Xaa at res.87 = (Ile, Val or Asn); Xaa at res.89 = (Lys or Arg); Xaa at res.90 = (Lys, Asn, Gln, His or Val); Xaa at res.91 = (Tyr or His); Xaa at res.92 = (Arg, Gln, Glu or Pro); Xaa at res.93 = (Asn, Glu or Asp); Xaa at res.95 = (Val, Thr, Ala or Ile); Xaa at res.97 = (Arg, Lys, Val, Asp or Glu); Xaa at res.98 = (Ala, Gly, Glu or Ser); Xaa at res.100 = (Gly or Ala); and Xaa at res.102 = (His or Arg).

[0033] Particularly useful sequences for use as morphogens include the C-terminal domains, e.g., the C-terminal 96-102 amino acid residues of Vgl, Vgr-1, DPP, OP-1, OP-2, CBMP-2A, CBMP-2B, GDF-1 (see Table II, below, and Seq. ID Nos. 5-14), as well as proteins comprising the C-terminal domains of 60A, BMP3, BMP5 and BMP6 (see Table II, below, and Seq. ID Nos. 24-28), all of which include at least the conserved six or seven cysteine skeleton. In addition, biosynthetic constructs designed from the generic sequences, such as COP-1, 3-5, 7, 16, disclosed in U.S. Pat. No. 5,011,691, also are useful. Other sequences include the inhibins/activin proteins (see, for example, U.S. Pat. Nos. 4,968,590 and 5,011,691). Accordingly, other useful sequences are those sharing at least 70% amino acid sequence homology or "similarity", and preferably 80% homology or similarity with any of the sequences above. These are anticipated to include allelic and species variants and mutants, and biosynthetic muteins, as well as novel members of this morphogenic family of proteins. Particularly envisioned in the family of related proteins are those proteins exhibiting morphogenic activity and wherein the amino acid changes from the preferred sequences include conservative changes, e.g., those as defined by Dayoff et al., *Atlas of Protein Sequence and Structure*; vol. 5, Suppl. 3, pp. 345-362, (M.O. Dayoff, ed., Nat'l BioMed. Research Fdn., Washington, D.C. 1979). As used herein, potentially useful sequences are aligned with a known morphogen sequence using the method of Needleman et al. ((1970) *J.Mol.Biol.* 48:443-453) and identities calculated by the Align program (DNASTar, Inc.). "Homology" or "similarity" as used herein includes allowed conservative changes as defined by Dayoff et al.

[0034] The currently most preferred protein sequences useful as morphogens include those having greater than 60% identity, preferably greater than 65% identity, with the amino acid sequence defining the conserved six cysteine skeleton of hOP1 (e.g., residues 43-139 of Seq. ID No.5). These most preferred sequences include both allelic and species variants of the OP-1 and OP-2 proteins, including the Drosophila 60A protein. Accordingly, useful morphogens include active proteins comprising species of polypeptide chains having the generic amino acid sequence herein referred to as "OPX", which accommodates the homologies between the various identified species of OP1 and OP2 (Seq. ID No. 29).

[0035] The morphogens useful in the use of this invention include proteins whether isolated from naturally-occurring sources, or produced by recombinant DNA or other synthetic techniques, and includes allelic and species variants of these proteins, naturally-occurring or biosynthetic mutants thereof, as well as various truncated and fusion constructs. Deletion or addition mutants also are envisioned to be active, including those which may alter the conserved C-terminal cysteine skeleton, provided that the alteration does not functionally disrupt the relationship of these cysteines in the folded structure. Accordingly, such active forms are considered the equivalent of the specifically described constructs disclosed herein. The proteins may include forms having varying glycosylation patterns, varying N-termini, a family of related proteins having regions of amino acid sequence homology, and active truncated or mutated forms of native or biosynthetic proteins, produced by expression of recombinant DNA in host cells.

[0036] The morphogenic proteins can be expressed from intact or truncated cDNA or from synthetic DNAs in prokaryotic or eucaryotic host cells, and purified, cleaved, refolded, and dimerized to form morphogenically active com-

positions. Currently preferred host cells include E. coli or mammalian cells, such as CHO, COS or BSC cells.

[0037] Thus, in view of this disclosure, skilled genetic engineers can isolate genes from cDNA or genomic libraries of various different species which encode appropriate amino acid sequences, or construct DNAs from oligonucleotides, and then can express them in various types of host cells, including both procaryotes and eucaryotes, to produce large quantities of active proteins capable of enhancing bone formation and/or inhibiting abnormal bone deterioration in a variety of mammals, including humans, for use in maintaining appropriate bone mass and bone remodeling in developing and adult bone tissue.

Brief Description of the Drawings

[0038] The foregoing and other objects and features of this invention, as well as the invention itself, may be more fully understood from the following description, when read together with the accompanying drawings, in which:

FIG. 1 compares the mitogenic effect of hOP-1 and TGF- β on rat osteoblasts;

FIG. 2 illustrates the effect of human osteogenic protein-1 (hOP-1) on the collagen synthesis of osteoblasts;

FIG. 3 compares the alkaline phosphatase induction effect of hOP-1 and TGF- β on rat osteoblasts;

FIG. 4 shows the long-term effect of hOP-1 on the production of alkaline phosphatase by rat osteoblasts;

FIG. 5 shows the effect of hOP-1 on parathyroid hormone-mediated cAMP production using rat osteoblasts in culture;

FIG. 6A and B graphs the effect of morphogen on osteoclastin synthesis (A), and the effect of morphogen on the rate of mineralization (B);

FIG. 7 shows Western Blot analysis of bovine colostrum using OP-1 and BMP2-specific antibodies;

FIG. 8A and B show results of in vivo and in vitro activity assays, respectively, for mammary extract purified OP-1;

FIG. 9 is a photomicrograph of an immunoblot showing the presence of hOP-1 in serum; and

FIG. 10 (A and B) are photomicrographs showing new endosteum bone formation following morphogen injection onto the endosteal surface (A), and new periosteum bone formation following morphogen injection onto the periosteal surface (B);

FIG. 11 is a graphic representation of the dose-dependent effect of morphogen on bone resorption; and

FIG. 12 (A and B) are schematic representations of morphogen inhibition of early mononuclear phagocytic cell multinuclearization in vivo;

Detailed Description of the Invention

[0039] It now has been discovered that the proteins described herein are effective agents for preventing loss of bone mass and/or for stimulating bone formation when provided systemically or injected directed into bone tissue in a mammal. As described herein, these proteins ("morphogens") may be used in the treatment of metabolic bone diseases and other disorders that cause an imbalance of the bone remodeling cycle, and/or which cause deterioration of the skeletal microstructure.

[0040] The invention is based on the discovery of a family of morphogenic proteins capable of inducing tissue morphogenesis in a mammal. More particularly, the invention is based on the discovery that these proteins play an important role, not only in embryogenesis, but also in the growth, maintenance and repair of bone tissue in juvenile and adult mammals.

[0041] It has been shown that implantation of a morphogen (including OP-1, CBMP2, DPP and 60A protein, and various biosynthetic constructs, such as COPS and COP7) together with a suitable matrix in subcutaneous sites in mammals induces a sequence of cellular events which leads to the formation of fully functional new bone, as determined by the specific activity of alkaline phosphatase, calcium content and histology of day 12 implants (see, for example, U.S. Pat. Nos. 4,968,590 and 5,011,691).

[0042] The morphogen-containing implants recruit nearby mesenchymal stem cells and trigger their differentiation into chondrocytes within 5-7 days. Upon capillary invasion, the chondrocytes hypertrophy, become calcified and subsequently are replaced by newly formed bone within 9-12 days. The mineralized bone then is remodeled extensively and becomes occupied by ossicles filled with functional bone marrow elements by 14-21 days.

[0043] As described herein, the morphogens provided herein stimulate the proliferation, growth and differentiation of osteoblasts in vitro (see Examples 2-7, below), and can induce bone formation in osteoporotic bone tissue in vivo when provided systemically to a mammal, or directly to bone tissue, without an associated matrix carrier (see Examples 8, 9, below.) In addition, the morphogens inhibit multinucleation of activated early mononuclear phagocytic cells (see Example 12, below). Moreover, inhibition of endogenous morphogen activity can inhibit normal skeleton development in a mammal (see Example 13, below.)

[0044] As described in Example 1 the naturally-occurring morphogens are widely distributed in the different tissues of the body. For example, as determined by northern blot hybridization, OP-1 is expressed primarily in the tissue of the urogenital tract (e.g., renal and bladder tissues). By contrast, Vgr-1, BMP3, BMP4 and BMP5 appear to be expressed primarily in the heart and lung. BMP5 also appears to be expressed significantly in liver tissue. GDF-1 appears to be expressed primarily in brain tissue. (See, for example, Ozkaynak et al. (1992) JBC, in publication.) Moreover, the tissue of synthesis may differ from the natural site of action of specific morphogens. For example, although OP-1 appears to be primarily synthesized in renal tissue, the protein is active in bone tissue. In addition, at least one morphogen, OP-1, is present in a number of body fluids, including saliva, milk (including mammary gland extract, colostrum and 57-day milk) and serum (see Example 11, below.) Accordingly, without being limited to a given theory, the morphogens described herein may behave as endocrine factors, e.g., proteins secreted from a factor-producing tissue in response to particular stimuli, and capable of being transported to, and acting on, a distant tissue. These findings further distinguish morphogens from other members of the TGF- β superfamily of proteins, including TGF- β , which act as local or autocrine factors produced by the tissue on which they act.

[0045] The pro domain may function to enhance protein solubility and/or to assist in tissue targeting of morphogens to particular tissues. For example, the mature, active form of OP-1 appears to be secreted from cells in association with the pro domain of the intact sequence. Accordingly, while, as explained herein, the morphogens useful in this invention have significant amino acid sequence homologies within the active domains and are similar in their ability to induce tissue morphogenesis, without being limited to any theory, it is hypothesized that the sequence variation within the morphogenic protein family members may reflect the different specific roles each morphogen plays in specific tissues under natural occurring conditions. For example, the significant sequence variation within the pro domains may mean that these regions of the protein sequence are important for targeting specific morphogens to different tissues for morphogenic activity therein.

[0046] Accordingly, the present disclosure comprises two fundamental aspects. In one aspect, the methods and compositions comprise a morphogen which, when administered to an individual, is capable of inhibiting loss of bone mass and/or stimulating bone formation in the individual. In another aspect, the methods and compositions comprise a morphogen-stimulating agent which, when administered to an individual, is capable of inducing the expression and/or secretion of sufficient endogenous morphogen within the individual to provide therapeutically effective concentrations capable of inhibiting loss of bone mass and/or stimulating bone formation in the individual.

[0047] Example 14 describes an assay for screening compounds to identify candidate morphogen-stimulating agents. Candidate agents then may be tested for their efficacy in vivo using, for example, the osteoporosis model described in Examples 8 and 9 below.

[0048] Provided below are detailed descriptions of suitable morphogens useful in the methods and compositions of this invention, as well as methods for the administration and application of these morphogens and/or of morphogen-stimulating agents. Also provided are numerous, nonlimiting examples which 1) illustrate the suitability of the morphogens and morphogen-stimulating agents described herein as therapeutic agents for inhibiting abnormal bone loss and/or for enhancing bone formation in a human, and 2) provide assays with which to test candidate morphogens and morphogen-stimulating agents for their efficacy.

I. Useful Morphogens

[0049] As defined herein a protein is morphogenic if it is capable of inducing the developmental cascade of cellular and molecular events that culminate in the formation of new, organ-specific tissue and comprises at least the conserved C-terminal six cysteine skeleton or its functional equivalent (see supra). Specifically, the morphogens generally are capable of all of the following biological functions in a morphogenically permissive environment: stimulating proliferation of progenitor cells; stimulating the differentiation of progenitor cells; stimulating the proliferation of differentiated cells; and supporting the growth and maintenance of differentiated cells, including the "redifferentiation" of transformed cells.

[0050] As disclosed therein, the morphogens may be purified from naturally-sourced material or recombinantly produced from procaryotic or eucaryotic host cells, using the genetic sequences disclosed therein. Alternatively, novel

10

15

40

50

10

15

20

30

35

40

45

50

55

10

15

20

25

30

35

40

45

50

55

10

15

20

30

35

40

45

50

55

	GDF-1	...	Ser	Pro	Phe	...
	60A	...	Gly	...	Leu	Pro	His
5	BMP5
	BMP6
					75					80
10	hOP-1	Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	Lys
	mOP-1
	hOP-2	...	Ser	...	Asn	Arg
15	mOP-2	...	Ser	...	Asn	Arg
	DPP	Asn	...	Gln	...	Thr	...	Val
	Vgl	...	Asn	Asn	Asp	Val	...	Arg
20	Vgr-1	Asn
	CBMP-2A	...	Glu	Asn	Glu	Lys	...	Val
	CBMP-2B	...	Glu	Tyr	Asp	Lys	...	Val
	BMP3	...	Glu	Asn	Lys	Val
25	GDF-1	...	Asn	...	Asp	Val	...	Arg
	60A	Leu	Asn	Asp	Glu	Asn
	BMP5
30	BMP6	Asn
						85				
35	hOP-1	Lys	Tyr	Arg	Asn	Met	Val	Val	Arg	
	mOP-1	
	hOP-2	...	His	Lys	
40	mOP-2	...	His	Lys	
	DPP	Asn	...	Gln	Glu	...	Thr	...	Val	
	Vgl	His	...	Glu	Ala	...	Asp	
	Vgr-1	
45	CBMP-2A	Asn	...	Gln	Asp	Glu	
	CBMP-2B	Asn	...	Gln	Glu	Glu	
	BMP3	Val	...	Pro	Thr	...	Glu	

	GDF-1	Gln	...	Glu	Asp	Asp
	60A	Ile	...	Lys
5	BMP5
	BMP6	Trp
		90					95		

10

	hOP-1	Ala	Cys	Gly	Cys	His
	mOP-1
15	hOP-2
	mOP-2
	DPP	Gly	Arg
20	Vgl	Glu	Arg
	Vgr-1
	CBMP-2A	Gly	Arg
	CBMP-2B	Gly	Arg
25	BMP3	Ser	...	Ala	...	Arg
	GDF-1	Glu	Arg
	60A	Ser
30	BMP5	Ser
	BMP6

100

**Between residues 56 and 57 of BMP3 is a Val residue;
 between residues 43 and 44 of GDF-1 lies
 the amino acid sequence Gly-Gly-Pro-Pro.

35

[0054] As is apparent from the foregoing amino acid sequence comparisons, significant amino acid changes can be made within the generic sequences while retaining the morphogenic activity. For example, while the GDF-1 protein sequence depicted in Table II shares only about 50% amino acid identity with the hOP-1 sequence described therein, the GDF-1 sequence shares greater than 70% amino acid sequence homology (or "similarity") with the hOP-1 sequence, where "homology" or "similarity" includes allowed conservative amino acid changes within the sequence as defined by Dayoff, et al., Atlas of Protein Sequence and Structure vol.5, supp.3, pp.345-362, (M.O. Dayhoff, ed., Nat'l BioMed. Res. Fd'n, Washington D.C. 1979.)

45

[0055] The currently most preferred protein sequences useful as morphogens in this disclosure include those having greater than 60% identity, preferably greater than 65% identity, with the amino acid sequence defining the conserved six cysteine skeleton of hOP-1 (e.g., residues 43-139 of Seq. ID No. 5). These most preferred sequences include both allelic and species variants of the OP-1 and OP-2 proteins, including the Drosophila 60A protein. Accordingly, the invention includes use of a morphogen in the manufacture of a medicament wherein the morphogen comprises a species of polypeptide chains having the generic amino acid sequence referred to herein as "OPX", which defines the seven cysteine skeleton and accommodates the identities between the various identified mouse and human OP1 and OP2 proteins. OPX is presented in Seq. ID No. 29. As described therein, each Xaa at a given position independently is selected from the residues occurring at the corresponding position in the C-terminal sequence of mouse or human OP1 or OP2 (see Seq. ID Nos. 5-8 and/or Seq. ID Nos. 16-23).

55

II. Formulations and Methods for Administering Therapeutic Agents

[0056] The morphogens may be provided to an individual by any suitable means, preferably directly, parenterally or orally. Where the morphogen is to be provided directly (e.g., locally, as by injection, to a bone tissue site), or parenterally, such as by intravenous, subcutaneous, intramuscular, intraorbital, ophthalmic, intraventricular, intracranial, intracapsular, intraspinal, intracisternal, intraperitoneal, buccal, rectal, vaginal, intranasal or by aerosol administration, the morphogen preferably comprises part of an aqueous solution. The solution is physiologically acceptable so that in addition to delivery of the desired morphogen to the patient, the solution does not otherwise adversely affect the patient's electrolyte and volume balance. The aqueous medium for the morphogen thus may comprise normal physiologic saline (9.85% NaCl, 0.15M), pH 7-7.4. The aqueous solution containing the morphogen can be made, for example, by dissolving the protein in 50% ethanol containing acetonitrile in 0.1% trifluoroacetic acid (TFA) or 0.1% HCl, or equivalent solvents. One volume of the resultant solution then is added, for example, to ten volumes of phosphate buffered saline (PBS), which further may include 0.1-0.2% human serum albumin (HSA). The resultant solution preferably is vortexed extensively. If desired, a given morphogen may be made more soluble by association with a suitable molecule. For example, association of the mature dimer with the pro domain of the morphogen increases solubility of the protein significantly. In fact, the endogenous protein is thought to be transported in this form. Another molecule capable of enhancing solubility and particularly useful for oral administrations, is casein. For example, addition of 0.2% casein increases solubility of the mature active form of OP-1 by 80%. Other components found in milk and/or various serum proteins also may be useful.

[0057] Useful solutions for oral or parenteral administration may be prepared by any of the methods well known in the pharmaceutical art, described, for example, in Remington's Pharmaceutical Sciences, (Gennaro, A., ed.), Mack Pub., 1990. Formulations may include, for example, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes, and the like. Formulations for direct administration, in particular, may include glycerol and other compositions of high viscosity. Biocompatible, preferably bioresorbable polymers, including, for example, hyaluronic acid, collagen, tricalcium phosphate, polybutyrate, lactide and lactide/glycolide copolymers, may be useful excipients to control the release of the morphogen in vivo. Other potentially useful parenteral delivery systems for these morphogens include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration contain as excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for parenteral administration may also include glycocholate for buccal administration, methoxysalicylate for rectal administration, or citric acid for vaginal administration.

[0058] Alternatively, the morphogens described herein may be administered orally. Oral administration of proteins as therapeutics generally is not practiced as most proteins readily are degraded by digestive enzymes and acids in the mammalian digestive system before they can be absorbed into the bloodstream. However, the morphogens described herein typically are acid-stable and protease-resistant (see, for example, U.S. Pat. No. 4,968,590.) In addition, at least one morphogen, OP-1, has been identified in bovine mammary gland extract, colostrum and milk (see Example 10, below) as well as saliva. Moreover, the OP-1 purified from mammary gland extract has been shown to be morphogenically active. Specifically, this protein has been shown to induce endochondral bone formation in mammals when implanted subcutaneously in association with a suitable matrix material, using a standard in vivo bone assay, such as is disclosed in U.S. Pat. No. 4,968,590. In addition, endogenous morphogen also has been detected in the bloodstream (see Example 11). These findings indicate that oral and parenteral administration are viable means for administering morphogens to an individual. In addition, while the mature forms of certain morphogens described herein typically are sparingly soluble, the morphogen form found in milk (and mammary gland extract and colostrum) is readily soluble, probably by association of the mature, morphogenically active form with the pro domain of the intact sequence and/or by association with one or more milk components. Accordingly, the compounds provided herein also may be associated with molecules capable of enhancing their solubility in vitro or in vivo, including, for example, part or all of a morphogen pro domain, and casein, as described above.

[0059] The compounds provided herein also may be associated with molecules capable of targeting the morphogen or morphogen-stimulating agent to bone tissue. For example, tetracycline and diphosphonates are known to bind to bone mineral, particularly at zones of bone remodeling, when they are provided systemically in a mammal. Alternatively, an antibody or other binding protein that interacts specifically with a surface molecule on bone tissue cells also may be used. Such targeting molecules further may be covalently associated to the morphogen or morphogen-stimulating agent with, for example, an acid labile bond such as an Asp-Pro linkage, using standard chemical means well known in the art. Because the local environment at bone remodeling sites is acidic, acid-labile linkages are expected to be preferentially cleaved at these sites, yielding active morphogen or morphogen-stimulating agent at the desired site. Useful targeting molecules may be designed, for example, using the single chain binding site technology disclosed, for example, in U.S. Pat. No. 5,091,513.

[0060] As described above, the morphogens provided herein share significant sequence homology in the C-terminal active domains. By contrast, the sequences diverge significantly in the sequences which define the pro domain. Accordingly, the pro domain may be morphogen-specific. As described above, it is also known that the various morphogens identified to date are differentially expressed in the different tissues. Accordingly, without being limited to any given theory, it is likely that, under natural conditions in the body, selected morphogens typically act on a given tissue. Accordingly, part or all of pro domains, which have been identified associated with the active form of the morphogen in solution, may serve as targeting molecules for the morphogens described herein. For example, the pro domains may interact specifically with one or more molecules at the target tissue to direct the morphogen associated with the pro domain to that tissue. Accordingly, another useful targeting molecule for targeting morphogen to bone tissue is part or all of a morphogen pro domain, particularly part or all of the pro domains of OP-1, BMP2 or BMP4, all of which proteins are found naturally associated with bone tissue.

[0061] Finally, the morphogens or morphogen-stimulating agents provided herein may be administered alone or in combination with other molecules known to have a beneficial effect on maintaining appropriate bone remodeling cycles in an individual at risk for excessive bone loss. Examples of useful cofactors include vitamin D₃, calcitonin, prostaglandins, parathyroid hormone, dexamethasone, estrogen and IGF.

[0062] The compounds provided herein can be formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable nontoxic excipients and carriers. As noted above, such compositions may be prepared for parenteral administration, particularly in the form of liquid solutions or suspensions; for oral administration, particularly in the form of tablets or capsules; or intranasally, particularly in the form of powders, nasal drops, or aerosols.

[0063] The compositions can be formulated for parenteral or oral administration to humans or other mammals in therapeutically effective amounts, e.g., amounts which provide appropriate concentrations of a morphogen to bone tissue for a time sufficient to inhibit loss of bone mass and/or to stimulate bone formation in individuals suffering from metabolic bone diseases and other bone remodeling disorders as described above. Therapeutic concentrations also are sufficient to repair fractures and other defects in skeletal microstructure, and to enhance maintenance of appropriate bone mass in developing juveniles and adults, including protecting individuals at risk for bone mass deterioration.

[0064] As will be appreciated by those skilled in the art, the concentration of the compounds described in a therapeutic composition will vary depending upon a number of factors, including the dosage of the drug to be administered, the chemical characteristics (e.g., hydrophobicity) of the compounds employed, and the route of administration. The preferred dosage of drug to be administered also is likely to depend on such variables as the type and extent of bone loss or defect, the overall health status of the particular patient, the relative biological efficacy of the compound selected, the formulation of the compound excipients, and its route of administration. In general terms, the compounds of this invention may be provided in an aqueous physiological buffer solution containing about 0.1 to 10% w/v compound for parenteral administration. Typical dose ranges are from about 10 ng/kg to about 1 g/kg of body weight per day; a preferred dose range is from about 0.1 µg/kg to 100 mg/kg of body weight per day. Optimally, the morphogen dosage given in all cases is between 2-20 µg of protein per kilogram weight of the patient per day. Currently preferred dose ranges for local injection of soluble morphogen to bone tissue are 0.1-50 µg morphogen/injection. No obvious morphogen-induced pathological lesions are induced when mature morphogen (e.g., OP-1, 20 µg) is administered daily to normal growing rats for 21 consecutive days. Moreover, 10 µg systemic injections of morphogen (e.g., OP-1) injected daily for 10 days into normal newborn mice does not produce any gross abnormalities.

III. Examples

Example 1. Identification of Morphogen-Expressing Tissue

[0065] Determining the tissue distribution of morphogens may be used to identify different morphogens expressed in a given tissue, as well as to identify new, related morphogens. Tissue distribution also may be used to identify useful morphogen-producing tissue for use in screening and identifying candidate morphogen-stimulating agents. The morphogens (or their mRNA transcripts) readily are identified in different tissues using standard methodologies and minor modifications thereof in tissues where expression may be low. For example, protein distribution may be determined using standard Western blot analysis or immunofluorescent techniques, and antibodies specific to the morphogen or morphogens of interest. Similarly, the distribution of morphogen transcripts may be determined using standard Northern hybridization protocols and transcript-specific probes.

[0066] Any probe capable of hybridizing specifically to a transcript, and distinguishing the transcript of interest from other, related transcripts may be used. Because the morphogens described herein share such high sequence homology in their active, C-terminal domains, the tissue distribution of a specific morphogen transcript may best be determined using a probe specific for the pro region of the immature protein and/or the N-terminal region of the mature protein. Another useful sequence is the 3' non-coding region flanking and immediately following the stop codon. These portions of the sequence vary substantially among the morphogens of this invention, and accordingly, are specific for each

protein. For example, a particularly useful Vgr-1-specific probe sequence is the PvuII-SacI fragment, a 265 bp fragment encoding both a portion of the untranslated pro region and the N-terminus of the mature sequence (see Lyons et al. (1989) PNAS 86:4554-4558 for a description of the cDNA sequence). Similarly, particularly useful mOP-1-specific probe sequences are the BstXI-BglI fragment, a 0.68 Kb sequence that covers approximately two-thirds of the mOP-1 pro region; a StuI-StuI fragment, a 0.2 Kb sequence immediately upstream of the 7-cysteine domain; and the EarI-PstI fragment, an 0.3 Kb fragment containing a portion of the 3'untranslated sequence (See Seq. ID No. 18, where the pro region is defined essentially by residues 30-291.) Similar approaches may be used, for example, with hOP-1 (Seq. ID No. 16) or human or mouse OP-2 (Seq. ID Nos. 20 and 22.)

[0067] Using these morphogen-specific probes, which may be synthetically engineered or obtained from cloned sequences, morphogen transcripts can be identified in mammalian tissue, using standard methodologies well known to those having ordinary skill in the art. Briefly, total RNA is prepared from various adult murine tissues (e.g., liver, kidney, testis, heart, brain, thymus and stomach) by a standard methodology such as by the method of Chomczynski et al. ((1987) Anal. Biochem 162:156-159) and described below. Poly (A)+ RNA is prepared by using oligo (dT)-cellulose chromatography (e.g., Type 7, from Pharmacia LKB Biotechnology, Inc.). Poly (A)+ RNA (generally 15 µg) from each tissue is fractionated on a 1% agarose/formaldehyde gel and transferred onto a Nytran membrane (Schleicher & Schuell). Following the transfer, the membrane is baked at 80°C and the RNA is cross-linked under UV light (generally 30 seconds at 1 mW/cm²). Prior to hybridization, the appropriate probe is denatured by heating. The hybridization is carried out in a lucite cylinder rotating in a roller bottle apparatus at approximately 1 rev/min for approximately 15 hours at 37°C using a hybridization mix of 40% formamide, 5 x Denhardt's, 5 x SSPE, and 0.1% SDS. Following hybridization, the non-specific counts are washed off the filters in 0.1 x SSPE, 0.1% SDS at 50°C.

[0068] Examples demonstrating the tissue distribution of various morphogens, including Vgr-1, OP-1, BMP2, BMP3, BMP4, BMP5, GDF-1, and OP-2 in developing and adult tissue are disclosed in co-pending USSN 752,764, and in Ozkaynak, et al., (1991) Biochem. Biophys. Res. Commn. 179:116-123, and Ozkaynak, et al. (1992) (JBC, in press), the disclosures of which are incorporated herein by reference. Using the general probing methodology described herein, northern blot hybridizations using probes specific for these morphogens to probe brain, spleen, lung, heart, liver and kidney tissue indicate that kidney-related tissue appears to be the primary expression source for OP-1, with brain, heart and lung tissues being secondary sources. Lung tissue appears to be the primary tissue expression source for Vgr-1, BMP5, BMP4 and BMP3. Lower levels of Vgr-1 also are seen in kidney and heart tissue, while the liver appears to be a secondary expression source for BMP5, and the spleen appears to be a secondary expression source for BMP4. GDF-1 appears to be expressed primarily in brain tissue. To date, OP-2 appears to be expressed primarily in early embryonic tissue. Specifically, northern blots of murine embryos and 6-day post-natal animals shows abundant OP2 expression in 8-day embryos. Expression is reduced significantly in 17-day embryos and is not detected in post-natal animals.

Example 2. Mitogenic Effect of Morphogen on Rat and Human Osteoblasts

[0069] The ability of a morphogen to induce proliferation of osteoblasts may be determined *in vitro* using the following assay. In this and all examples involving osteoblast cultures, rat osteoblast-enriched primary cultures preferably are used. Although these cultures are heterogeneous in that the individual cells are at different stages of differentiation, the culture is believed to more accurately reflect the metabolism and function of osteoblasts *in vivo* than osteoblast culture obtained from established cell lines. Unless otherwise indicated, all chemicals referenced are standard, commercially available reagents, readily available from a number of sources, including Sigma Chemical, Co., St. Louis; Calbiochem, Corp., San Diego, and Aldrich Chemical Co., Milwaukee.

[0070] Rat osteoblast-enriched primary cultures were prepared by sequential collagenase digestion of newborn suture-free rat calvaria (e.g., from 1-2 day-old animals, Long-Evans strain, Charles River Laboratories, Wilmington, MA), following standard procedures, such as are described, for example, in Wong et al., (1975) PNAS 72:3167-3171. Rat osteoblast single cell suspensions then were plated onto a multi-well plate (e.g., a 48 well plate) at a concentration of 50,000 osteoblasts per well in alpha MEM (modified Eagle's medium, Gibco, Inc., Long Island) containing 10% FBS (fetal bovine serum), L-glutamine and penicillin/streptomycin. The cells were incubated for 24 hours at 37°C, at which time the growth medium was replaced with alpha MEM containing 1% FBS and the cells incubated for an additional 24 hours so that cells were in serum-deprived growth medium at the time of the experiment.

[0071] The cell culture then was divided into three groups: (1) wells which received 0.1, 1.0, 10.0, 40 and 80.0 ng of morphogen; (2) wells which received 0.1, 1.0, 10.0 and 40 ng of a local-acting growth factor; and (3) the control group, which received no growth factors. In this example, OP-1 was the morphogen tested, and TGF-β was the local-acting growth factor. The cells then were incubated for an additional 18 hours after which the wells were pulsed with 2µCi/well of ³H-thymidine and incubated for six more hours. The excess label then was washed off with a cold solution of 0.15 M NaCl, 250 µl of 10% trichloroacetic acid then was added to each well and the wells incubated at room temperature for 30 minutes. The cells then were washed three times with cold distilled water, and lysed by the addition of 250

μl of 1% sodium dodecyl sulfate (SDS) for a period of 30 minutes at 37°C. The cell lysates then were harvested using standard means well known in the art, and the incorporation of ³H-thymidine into cellular DNA was determined by liquid scintillation as an indication of mitogenic activity of the cells. The results, shown in FIG. 1, demonstrate that OP-1 (identified in the figure by squares) stimulates ³H-thymidine incorporation into DNA, and thus promotes osteoblast cell proliferation. The mitogenesis stimulated by 40 ng of OP-1 in serum-free medium was equivalent to the mitogenic effect of 10% fresh serum alone. By contrast, the effect of TGF-β (indicated by diamonds in Fig. 1) is transient and biphasic. At high concentrations, TGF-β has no significant effect on osteoblast cell proliferation. This system may be used to test other morphogens for their effect on cell proliferation.

[0072] The in vitro effect of a morphogen on osteoblast proliferation also was tested on human primary osteoblasts (obtained from bone tissue of a normal adult patient and prepared as described above) and on osteosarcoma-derived cells, and in all cases induced cell proliferation. In addition, similar experiments, performed using BMP4 (BMP2B) and BMP3 shows these morphogens also can stimulate osteoblast proliferation and growth. (See Chen et al., (1991) J. Bone and Min. Res. 6: 1387-1393, and Vukicevic, (1989) PNAS 86: 8793-8797.)

[0073] The effect of a given morphogen on bone cell growth and/or development also may be tested using a variety of bone cell markers: e.g., collagen synthesis, alkaline phosphatase activity, parathyroid hormone-mediated cyclic AMP (cAMP) production, osteocalcin synthesis, and by assessing the rate of mineralization in osteoblasts. Of these, alkaline phosphatase activity, parathyroid hormone-mediated cAMP production, osteocalcin synthesis and mineralization promotion are specific markers for the differentiated osteoblast phenotype. Experimental systems for testing these parameters as well as collagen synthesis are described below in Examples 3-7. In all cases morphogen alone stimulated expression of these phenotype-specific markers. In Examples 3-7 OP-1 was the morphogen tested. Similar experiments, performed using BMP4 (BMP2B) shows that this morphogen also induces osteoblast differentiation. (See Chen, et al. (1991) T. Bone and Min. Res. 6: 1387-1392, and Vukicevic, (1989) PNAS 86: 8793-8797.)

Example 3. Effect of Morphogen on Collagen Synthesis in Rat Osteoblasts

[0074] The effect of a morphogen on collagen production in rat osteoblasts in vitro may be determined as follows.

[0075] Rat osteoblasts were prepared and cultured in a multi-well plate as described for Example 2. In this example a 24-well plate was used. The cultured cells then were divided into three groups: (1) wells which received 1, 10 or 40 ng of morphogen per ml of medium; (2) wells which received 1, 10 or 40 ng of a local-acting growth factor per ml of medium; and (3) a control group which received no growth factors. In this example, OP-1 was the morphogen tested, and TGF-β was the local-acting growth factor.

[0076] The samples were incubated for 68 hours at 37°C with 5% CO₂ in a humidified incubator. Twenty-five (25) μCi of ³H proline were added into each well and incubated for six additional hours. The cells then were frozen at -20°C until the collagen assay was performed. The cells then were assayed for collagen production by detecting incorporation of ³H-proline into total collagenase-digestible protein (CDP). The results, shown in FIG. 2, demonstrate that OP-1 stimulates type-I collagen synthesis, as measured by ³H-proline incorporation into total CDP. Thus, OP-1 promotes collagen synthesis in vitro by preosteoblasts and mature osteoblasts.

Example 4. Alkaline Phosphatase Induction of Osteoblasts by Morphogen

4.1 Morphogen-specific Alkaline Phosphatase Induction

[0077] Since alkaline phosphatase production is an indicator of bone formation by differentiated, functional osteoblasts, a morphogen may be assessed for its potential osteogenic effects using this osteoblast marker in the following in vitro test system.

[0078] Rat osteoblasts were prepared and cultured in a multi-well plate as described for Example 2. In this example a 24-well plate was used. The cultured cells then were divided into three groups: (1) wells which received varying concentrations of morphogen; (2) wells which received varying concentrations of a local-acting growth factor; and (3) a control group which received no growth factors. In this example OP-1 was the morphogen tested at the following concentrations: 0.1, 1.0, 10.0, 40.0 or 80.0 ng/ml medium; and TGF-β was the local-acting growth factor, tested at 0.1, 1.0, 10.0, 40.0 or 80.0 ng/ml medium. The cells then were incubated for 72 hours. After the incubation period the cell layer was extracted with 0.5 ml of 1% Triton X-100. The resultant cell extract was centrifuged, 100 μl of the extract was added to 90 μl of paranitrosophenylphosphate (PNPP)/glycerine mixture and incubated for 30 minutes in a 37°C water bath and the reaction stopped with 100 μl NaOH. The samples then were run through a plate reader (e.g., Dynatech MR700 plate reader, and absorbance measured at 400 nm, using p-nitrophenol as a standard) to determine the presence and amount of alkaline phosphate activity. Protein concentrations were determined by the Biorad method. Alkaline phosphatase activity was calculated in units/μg protein, where 1 unit=1 nmol p-nitrophenol liberated/30 minutes at 37°C.

[0079] The results, shown in FIG. 3, illustrate that morphogen alone stimulates the production of alkaline phosphatase

in osteoblasts, and thus promotes the growth and expression of the osteoblast differentiated phenotype. In the figure, squares represent OP-1 concentrations, and diamonds represent TGF- β concentrations.

4.2. Long Term Effect of Morphogen on the Production of Alkaline Phosphatase by Rat Osteoblasts

[0080] In order to determine the long term effect of a morphogen on the production of alkaline phosphatase by rat osteoblasts, the following assay may be performed.

[0081] Rat osteoblasts were prepared and cultured in multi-well plates as described in Example 2. In this example six sets of 24 well plates are plated with 50,000 rat osteoblasts per well. The wells in each plate, prepared as described above, then were divided into three groups: (1) those which received 1 ng of morphogen per ml of medium; (2) those which received 40 ng of morphogen/ml of medium; and (3) those which received 80 ng of morphogen/ml of medium. Each plate then was incubated for different lengths of time: 0 hours (control time), 24 hours, 48 hours, 96 hours, 120 hours and 144 hours. After each incubation period, the cell layer was extracted with 0.5 ml of 1% Triton X-100. The resultant cell extract was centrifuged, and alkaline phosphatase activity determined as for Example 4, using paranitrosophenylphosphate (PNPP). The results, shown in FIG. 4, illustrate that morphogen alone stimulates the production of alkaline phosphatase in osteoblasts, that increasing doses of OP-1 further increase the level of alkaline phosphatase production, and that the morphogen-stimulated elevated levels of alkaline phosphatase in the treated osteoblasts lasts for an extended period of time. In the figure, circles represent 1 ng OP-1; squares, 40 ng OP-1; and diamonds, 80 ng OP-1.

Example 5. Morphogen-Induced Parathyroid Hormone Mediated cAMP Production in Rat Osteoblasts

[0082] The effect of a morphogen on parathyroid hormone-mediated cAMP production in rat osteoblasts in vitro may be determined as follows.

[0083] Rat osteoblasts were prepared and cultured in a multiwell plate as described for Example 2 above. In this example a 24-well plate was used. The cultured cells then were divided into three groups: (1) wells which received varying concentrations of morphogen (in this example, OP-1, at 1.0, 10.0 and 40.0 ng/ml medium); (2) wells which received varying concentrations of a local-acting growth factor (in this example, TGF- β , at 0.1, 1.0, and 5.0 ng/ml medium); and (3) a control group which received no growth factors. The plate was then incubated for another 72 hours. At the end of the 72 hours the cells were treated with medium containing 0.5% bovine serum albumin (BSA) and 1mM 3-isobutyl-1-methyl xanthine for 20 minutes followed by the addition into half of the wells of human recombinant parathyroid hormone (hPTH, Sigma, St. Louis) at a concentration of 200ng/ml for 10 minutes. The cell layer was extracted from each well with 0.5 ml of 1% Triton X-100. The cAMP levels were then determined using a radioimmunoassay kit (Amersham, Arlington Heights, Illinois). The results, shown in FIG. 5, demonstrate that morphogen alone stimulates an increase in the PTH-mediated cAMP response, and thus promotes the growth and expression of the osteoblast differentiated phenotype.

Example 6. Effect of Morphogen on Osteocalcin Synthesis and the Rate of Mineralization by Osteoblasts in Culture

[0084] Osteocalcin is a bone-specific protein synthesized by osteoblasts which plays an integral role in the rate of bone mineralization in vivo. Circulating levels of osteocalcin in serum are used as a marker for osteoblast activity and bone formation in vivo. Induction of osteocalcin synthesis in osteoblast-enriched cultures can be used to assay morphogen efficacy in vitro.

[0085] Rat osteoblasts are prepared and cultured in a multi-well plate as for Example 2. In this example cells were cultured in a 24-well plate. In this experiment the medium was supplemented with 10%FBS, and on day 2, cells were fed with fresh medium supplemented with fresh 10 mM β -glycerophosphate (Sigma, Inc.). Beginning on day 5 and twice weekly thereafter, cells were fed with a complete mineralization medium containing all of the above components plus fresh L(+)-ascorbate, at a final concentration of 50 μ g/ml medium. Morphogen then was added to the wells directly. In this example, OP-1 in 50% acetonitrile (or 50% ethanol) containing 0.1% trifluoroacetic acid (TFA) was added at no more than 5 μ l morphogen/ml medium. Control wells received solvent vehicle only. The cells then were re-fed and the conditioned medium sample diluted 1:1 in standard radioimmunoassay buffer containing standard protease inhibitors and stored at -20° C until assayed for osteocalcin. Osteocalcin synthesis then was measured by standard radioimmunoassay using a commercially available rat osteocalcin-specific antibody.

[0086] Mineralization was determined on long term cultures (13 day) using a modified von Kossa staining technique on fixed cell layers: cells were fixed in fresh 4% paraformaldehyde at 23° C for 10 mn, following rinsing cold 0.9% NaCl. Fixed cells then were stained for endogenous alkaline phosphatase at pH 9.5 for 10 min, using a commercially available kit (Sigma, Inc.) Purple stained cells then were dehydrated with methanol and air dried after 30 min incubation in 3% AgNO₃ in the dark, H₂O-rinsed samples were exposed for 30 sec to 254 nm UV light to develop the black silver-

stained phosphate nodules. Individual mineralized foci (at least 20 μm in size) were counted under a dissecting microscope and expressed as nodules/culture (see Fig. 6B).

[0087] As can be seen in Fig. 6A OP-1 stimulates osteocalcin synthesis in osteoblast cultures. The increased osteocalcin synthesis in response to OP-1 is dose dependent and showed a 5-fold increase over the basal level using 25 ng of OP-1/10 ml medium after 13 days of incubation. The enhanced osteocalcin synthesis also can be confirmed by detecting the elevated osteocalcin mRNA message (20-fold increase) using a rat osteocalcin-specific probe. In addition, the increase in osteocalcin synthesis correlates with increased mineralization in long term osteoblast cultures as determined by the appearance of mineral nodules (compare Fig. 6A and 6B.) OP-1 increases the initial mineralization rate about 20-fold compared to untreated cultures. Similar experiments performed using TGF- β indicate that TGF- β does not induce osteocalcin synthesis or promote the mineralization process. Thus, morphogen alone promotes the growth and expression of the osteoblast differentiated phenotype.

Example 7. Effect of Morphogen on Bone Derived Growth Factor Induction in vitro

[0088] IGF-I and IGF-II are bone-derived growth factors involved in coupling bone formation with bone resorption in the bone remodeling cycle. The effect of morphogen on the production of these and other bone-derived growth factors, including TGF- β , may be evaluated using the following procedure.

[0089] Rat or human osteoblasts were prepared and cultured in a multiwell plate as for Example 2. The wells of the plate were divided in to groups in which different concentrations of morphogen were added (e.g., 0, 1, 10, and 100 ng). In this example, OP-1 was the morphogen used. The plate then was incubated for a prescribed period of time, e.g., 72 hours, and the level of IGF detected, e.g., by immunolocalization, using a commercially available antibody specific for IGFs. OP-1 induced the level of both IGF-I and IGF-II significantly. Greater than six fold IGF-I and two fold IGF-II were induced following exposure to 100 ng OP-1/ml. In addition, OP-1 stimulated production of the IGF-I stimulating factor, BP3 (IGF-I binding protein 3).

Example 8. Effect of Morphogen on Trabecular Bone in Ovariectomized (OVX) Rats

[0090] As indicated above, serum alkaline phosphatase and osteocalcin levels are indicators of bone formation within an individual. In order to determine the effect of a morphogen on bone production *in vivo*, these parameters are measured under conditions which promote osteoporosis, e.g., wherein osteoporosis is induced by ovary removal in rats.

[0091] Forty Long-Evans rats (Charles River Laboratories, Wilmington) weighing about 200g each are ovariectomized (OVX) using standard surgical procedures, and ten rats are sham-operated. The ovariectomization of the rats produces an osteoporotic condition within the rats as a result of decreased estrogen production. Food and water are provided *ad libitum*. Eight days after ovariectomy, the rats, prepared as described above, were divided into five groups: (A), 10 sham-operated rats; (B), 10 ovariectomized rats receiving 1 ml of phosphate-buffered saline (PBS) i.v. in the tail vein; (C) 10 ovariectomized rats receiving about 1 mg of $17\beta\text{E}_2$ (17β -estradiol E_2) by intravenous injection through the tail vein; (D) 9 ovariectomized rats receiving daily injections of approximately $2\mu\text{g}$ of morphogen by tail vein for 22 days; and (E) 9 ovariectomized rats receiving daily injections of approximately $20\mu\text{g}$ of morphogen by tail vein for 22 days. In this example, OP-1 was the morphogen tested.

[0092] On the 15th and 21st day of the study, each rat was injected with 5 mg of tetracycline, and on day 22, the rats were sacrificed. The body weights, uterine weights, serum alkaline phosphate levels, serum calcium levels and serum osteocalcin levels then were determined for each rat. The results are shown in Tables III and IV.

Table III

Body Weights, Uterine Weights and Alkaline Phosphatase			
Group	Body Weights	Uterine Weights	Alk. Phosphatase
	(g)	(g)	(U/L)
A-SHAM	250.90 \pm 17.04	0.4192 \pm 0.10	43.25 \pm 6.11
B-OVX+PBS	273.40 \pm 16.81	0.1650 \pm 0.04	56.22 \pm 6.21
C-OVX+E2	241.66 \pm 21.54	0.3081 \pm 0.03	62.66 \pm 4.11

Table III (continued)

Body Weights, Uterine Weights and Alkaline Phosphatase			
Group	Body Weights	Uterine Weights	Alk. Phosphatase
	(g)	(g)	(U/L)
D-OVX+OP-1 (2 μ g)	266.67 \pm 10.43	0.1416 \pm 0.03	58.09 \pm 12.97
E-OVX+OP-1 (20 μ g)	272.40 \pm 20.48	0.1481 \pm 0.05	66.24 \pm 15.74

TABLE IV

Serum Calcium and Serum Osteocalcin Levels		
Group	Serum Calcium	Serum Osteocalcin
	(ng/dl)	(ng/ml)
A-SHAM	8.82 \pm 1.65	64.66 \pm 14.77
B-OVX+PBS	8.95 \pm 1.25	69.01 \pm 10.20
C-OVX+E2	9.20 \pm 1.39	67.13 \pm 17.33
D-OVX+OP-1 (2 μ g)	8.77 \pm 0.95	148.50 \pm 84.11
E-OVX+OP-1 (20 μ g)	8.67 \pm 1.94	182.42 \pm 52.11

[0093] The results presented in Table III and IV show that intravenous injection of morphogen into ovariectomized rats produces a significant increase in serum alkaline phosphatase and serum osteocalcin levels and demonstrates that systemic administration of the morphogen stimulates bone formation in osteoporotic bone.

Example 9. Histomorphometric Analysis of Morphogen on the Tibia Diaphysis in Ovariectomized (OVX) Rats

[0094] Fifteen female Long-Evans rats weighing about 160 g were ovariectomized (OVX) to produce an osteoporotic condition and five rats were sham operated (Charles River Laboratories, Wilmington, MA.) as described for Example 8. Food and water were provided ad libitum. Twenty-two days after ovariectomy, the rats were divided into four groups: (A) sham-operated (1 ml of PBS by intravenous injection through tail vein (5 rats); (B) OVX, into which nothing was injected (5 rats); (C) OVX, receiving about 1 mg of 17 β E₂ by intravenous injection through the tail vein (5 rats), and (D) OVX, receiving about 1 μ g of morphogen by intravenous injection through the tail vein (5 rats). In this example, OP-1 was morphogen tested.

[0095] The rats were injected daily as described for seven days, except no injections were given on the thirteenth day. The rats then were sacrificed on the nineteenth day. The tibial diaphyseal long bones then were removed and fixed in ethanol and histomorphometric analysis was carried out using standard procedures well known in the art. The results are shown in Table V.

Table V

	(A)	(B)	(C)	(D)
MEASUREMENT	CONTROL	OVX	OVX + E ₂	OVX + OP-1
Longitudinal Growth Rate (μ m/day)	20.2 \pm 0.3	19.4 \pm 0.2	4.9 \pm 0.5	17.9 \pm 0.9
Cancellous Bone Volume (BV/TV, bone vol/total vol)	20.2 \pm 1.5	13.0 \pm 1.6	13.7 \pm 2.1	16.6 \pm 1.8
Cancellous Bone Perimeter (mm)	16.2 \pm 1.8	9.6 \pm 0.9	11.5 \pm 1.1	12.2 \pm 0.7
Labeled Cancellous Perimeter (%)	35.5 \pm 1.5	51.9 \pm 5.6	58.0 \pm 4.2	39.2 \pm 1.9
Mineral Apposition Rate (μ m/day)	1.76 \pm 0.14	2.25 \pm 0.16	1.87 \pm 0.08	1.86 \pm 0.20

[0096] The results presented in Table V confirm the results of Example 8, that intravenous injection of OP-1 into ovariectomized rats stimulates bone growth for bone which had been lost due to the drop in estrogen within the individual

rat. Specifically, the inhibition of cancellous bone volume in OVX rats is repaired by the systemically provided morphogen. In addition, in morphogen-treated rats the labelled cancellous perimeter and mineral apposition rate now return to levels measured in the control, sham-operated rats. Moreover, morphogen treatment does not inhibit longitudinal bone growth, unlike estrogen treatment, which appears to inhibit bone growth significantly. Accordingly, systemic administration of a morphogen in therapeutically effective concentrations effectively inhibits loss of bone mass in a mammal without inhibiting natural bone formation.

Example 10. Determination of the Presence of Morphogen in Body Fluids

[0097] OP-1 has been identified in saliva, human blood serum, and various milk forms, including mammary gland extract, colostrum, and 57-day bovine milk. Moreover, as described below, the body fluid extracted protein is morphogenically active. The discovery that the morphogen naturally is present in milk, together with the known observation that mature, active OP-1 is acid-stable and protease-resistant, indicate that oral administration is a useful route for therapeutic administration of morphogen to a mammal. Oral administration typically is the preferred mode of delivery for extended or prophylactic therapies. In addition, the identification of morphogen in all milk forms, including colostrum, indicates that the protein plays a significant role in tissue development, including skeletal development of juveniles (see Example 13, below).

10.1 Morphogen Detection in Milk

[0098] OP-1 was partially purified from rat mammary gland extract and bovine colostrum and 57 day milk by passing these fluids over a series of chromatography columns: (e.g., cation-exchange, affinity and reverse phase). At each step the eluant was collected in fractions and these were tested for the presence of OP-1 by standard immunoblot. Immunoreactive fractions then were combined and purified further. The final, partially purified product then was examined for the presence of OP-1 by Western blot analysis using OP-1-specific antisera, and tested for in vivo and in vitro activity.

[0099] OP-1 purified from the different milk sources were characterized by Western blotting using antibodies raised against OP-1 and BMP2. Antibodies were prepared using standard immunology protocols well known in the art, and as described generally in Example 14, below, using full-length *E. coli*-produced OP-1 and BMP2 as the immunogens.

[0100] As shown in Fig. 7 OP-1 purified from colostrum reacts with the anti-OP-1 antibody, but not with anti-BMP2 antibody. In Fig. 7 lane 1 contains reduced, purified, recombinantly-produced OP-1; lane 2 contains purified bovine colostrum, and lane 3 contains reduced COP-16, a biosynthetic construct having morphogenic activity and an amino acid sequence modeled on the proteins described herein, but having highest amino acid sequence homology with BMP2 (see US Pat. No. 5,011,691 for the COP-16 amino acid sequence.) In Fig. 7A the gel was probed with anti-OP-1 antibody; in Fig. 7B, the gel was probed with anti-BMP2 antibody. As can be seen in the figure, anti-OP-1 antibody hybridizes only with protein in lanes 1 and 2, but not 3; while anti-BMP2 antibody hybridizes with lane 3 only.

[0101] Column-purified mammary gland extract and 57-day milk also reacts specifically with anti-OP-1 antibodies, including antibody raised against the full length *E. coli* OP-1, full length mammalian-produced OP-1, and the OP-1 Ser-17-Cys peptide (e.g., the OP-1 N-terminal 17 amino acids).

[0102] The morphogenic activity of OP-1 purified from mammary gland extract was evaluated in vivo as follows. A sample was prepared from each OP-1 immunoreactive fraction of the mammary gland extract-derived OP-1 final product by lyophilizing a portion (33%) of the fraction and resuspending the protein in 220 μ l of 50% acetonitrile/0.1% TFA. After vortexing, 25 mg of collagen matrix was added. The samples were lyophilized overnight, and implanted in Long Evans rats (Charles River Laboratories, Wilmington, MA, 28-35 days old). Each fraction was implanted in duplicate. For details of the collagen matrix implantation procedure, see, for example, U.S. Pat. No. 4,968,590, hereby incorporated by reference. After 12 days, the implants were removed and evaluated for new bone formation by histological observation.

[0103] The results are presented in Fig.8A, where "% activity" refers to the percent of bone formation/total area covered by bone in the histology sample. In the figure, solid bars represent implants using mammary extract-derived OP-1, each bar corresponding to an immunoreactive fraction of the purified product, the fraction number being indicated on the x-axis. The hatched bar represents an implant using recombinantly produced OP-1 (600 ng). As can be seen in the figure, all immunoreactive fractions are osteogenically active.

[0104] Similarly, the morphogenic activity of OP-1 purified from mammary gland extract was evaluated in vitro by measuring alkaline phosphatase activity in vitro using the following assay. Test samples were prepared as for the in vivo assay, using 15-20% of individual immunoreactive fractions collected from the final product. Alkaline phosphatase activity was tested as described above in Example 4. The results, presented in Fig. 8B, indicate that the immunoreactive fractions can stimulate alkaline phosphatase activity in vitro. Moreover, the activity correlates well with that produced by highly purified, recombinantly produced, OP-1. In Fig. 8B solid bars represent assays performed with mammary

gland-purified OP-1, each bar corresponding to an immunoreactive fraction of column-purified OP-1, the fraction numbers being indicated on the x-axis; the hatched bar represents the assay performed with purified, recombinantly-produced OP-1 (100 ng/ml); and the cross-hatched bar represents background.

5 10.2 Morphogen Detection in Serum

[0105] Morphogen may be detected in serum using morphogen-specific antibodies. The assay may be performed using any standard immunoassay, such as Western blot (immunoblot) and the like. Preferably, the assay is performed using an affinity column to which the morphogen-specific antibody is bound and through which the sample serum then is poured, to selectively extract the morphogen of interest. The morphogen then is eluted. A suitable elution buffer may be determined empirically by determining appropriate binding and elution conditions first with a control (e.g., purified, recombinantly-produced morphogen.) Fractions then are tested for the presence of the morphogen by standard immunoblot, and the results confirmed by N-terminal sequencing. Preferably, the affinity column is prepared using monoclonal antibodies. Morphogen concentrations in serum or other fluid samples then may be determined using standard protein quantification techniques, including by spectrophotometric absorbance or by quantitation of conjugated antibody.

[0106] Presented below is a sample protocol for identifying OP-1 in serum. Following this general methodology other morphogens may be detected in body fluids, including serum. The identification of morphogen in serum further indicates that systemic administration is a suitable means for providing therapeutic concentrations of a morphogen to an individual, and that morphogens likely behave systemically as endocrine-like factors. Finally, using this protocol, fluctuations in endogenous morphogen levels can be detected, and these altered levels may be used as an indicator of bone tissue dysfunction. Alternatively, fluctuations in morphogen levels may be assessed by monitoring morphogen transcription levels, either by standard northern blot analysis as described in Example 1, or by *in situ* hybridization, using a labelled probe capable of hybridizing specifically to morphogen RNA, and standard RNA hybridization protocols well described in the art and described generally in Example 1.

[0107] OP-1 was detected in human serum using the following assay. A monoclonal antibody raised against mammalian, recombinantly produced OP-1 using standard immunology techniques well described in the art and described generally in Example 14, was immobilized by passing the antibody over an agarose-activated gel (e.g., Affi-Gel™, from Bio-Rad Laboratories, Richmond, CA, prepared following manufacturer's instructions) and used to purify OP-1 from serum. Human serum then was passed over the column and eluted with 3M K-thiocyanate. K-thiocyanate fractions then were dialyzed in 6M urea, 20mM PO₄, pH 7.0, applied to a C8 HPLC column, and eluted with a 20 minute, 25-50% acetonitrile/0.1% TFA gradient. Mature, recombinantly produced OP-1 homodimers elute between 20-22 minutes. Fractions then were collected and tested for the presence of OP-1 by standard immunoblot using an OP-1 specific antibody as for Example 10.A. Fig. 9 is an immunoblot showing OP-1 in human sera under reducing and oxidized conditions. In the figure, lanes 1 and 4 are OP-1 standards, run under oxidized (lane 1) and reduced (lane 4) conditions. Lane 5 shows molecular weight markers at 17, 27 and 39 kDa. Lanes 2 and 3 are human sera OP-1, run under oxidized (lane 2) and reduced (lane 3) conditions.

[0108] Morphogens may be used in diagnostic applications by comparing the quantity of morphogen present in a body fluid sample with a predetermined reference value, with fluctuations in fluid morphogen levels indicating a change in the status of bone tissue. Alternatively, fluctuations in the level of endogenous morphogen antibodies may be detected by this method, most likely in serum, using an antibody or other binding protein capable of interacting specifically with the endogenous morphogen antibody. Detected fluctuations in the levels of the endogenous antibody may be used as indicators of a change in tissue status.

45 Example 11. Morphogen-induced Periosteal and Endosteal Bone Formation

[0109] Osteoclast-induced bone resorption occurs primarily at the endosteal surface of bone tissue. Accordingly, in bone remodeling disorders the marrow cavity is enlarged unnaturally, weakening the weight bearing capacity of the remaining bone. The following example provides means for evaluating the ability of the morphogens described herein to increase endosteal and periosteal bone mass in a mammal. In this example, both periosteal and endosteal bone formation are induced by direct injection of a morphogen in a biocompatible solution directly to the bone tissue. As demonstrated below, morphogens can induce new bone formation and increase bone mass at both surfaces when provided to the bone by direct injection. Direct injection may be a preferred mode of administration for providing therapeutically effective concentrations to reduce an enlarged marrow cavity, and/or to repair fractures and other damage to bone tissue microstructure.

[0110] Morphogen was provided to either the periosteum (outer or peripheral bone surface) and endosteum (interior bone surface, e.g., that surface lining the marrow cavity) of a rat femur by a single injection in each case. Specifically, morphogen (e.g., OP-1, 2-20 µg) was provided to the bone tissue as an insoluble colloidal suspension in phosphate-

buffered saline. Endosteal injection was performed through a microhole made with a hand-held orthopedic drill. After 7 days, the treated bones were removed and prepared for histological evaluation as described in U.S. Pat. No. 4,968,590. As little as 2 µg morphogen is sufficient to induce new bone formation at the site of injection within 4-7 days. In addition, bone induction is dose-dependent. Photomicrographs of the histology are presented in Fig. 10. In the figure, "ob" means old bone, "bm" means bone marrow, "nb" means new bone, and "m" means muscle. Fig. 10A shows new bone formed following injection of morphogen to the endosteal surface. As can be seen in the figure, new bone has formed within the bone marrow cavity, filling in the periphery of the cavity. Fig 10B shows new bone formed following injection of morphogen to the periosteal surface, replacing the muscle normally present.

Example 12. Effect of Morphogen on Bone Resorption

[0111] The effect of morphogen on bone resorption may be evaluated using rat osteoclasts on bovine bone slices, in the presence and absence of morphogen, and the effect of morphogen on pit formation (resorption index) determined. Under standard conditions rat osteoclasts begin resorbing the bone tissue, causing pit formation on the bone slice surface. In this experiment OP-1 was the morphogen tested, at concentrations of 0, 5, 10, 20, 40, 50, and 100 ng/ml.

[0112] The results are presented in figure 11, where the resorption index is calculated as a percent of the control (e.g., bone resorption in the absence of morphogen), calculated as the number of pits per a given slice surface area. Below 40 ng bone resorption is enhanced; above 40 ng, OP-1 has no apparent effect on bone resorption. The results highlight the integral role the morphogen plays in bone remodeling. OP-1 is stored in bone tissue *in vivo*. In a normal bone remodeling cycle, the local concentration of OP-1 at the surface likely is low when osteoclasts begin resorbing bone, and the low concentration may enhance and/or stimulate bone resorption. As resorption continues, the local concentration of OP-1 at the surface likely increases, to a concentration that no longer has an effect on osteoclasts, but continues to affect osteoblast growth and activity (see Examples 2-7), stimulating bone growth.

[0113] In addition, morphogens can inhibit multinucleation of mononuclear phagocytic cells under conditions where these cells normally would be activated. For example, in the absence of morphogen, an implanted substrate material (e.g., implanted subcutaneously) composed of, for example, mineralized bone, a ceramic such as titanium oxide or any other substrate that provokes multinucleated giant cell formation, rapidly becomes surrounded by multinucleated giant cells, e.g., activated phagocytes stimulated to respond and destroy the foreign object. In the presence of morphogen however, the recruited cells remain in their mononuclear precursor form and the matrix material is undisturbed. Figure 12 illustrates this effect of morphogens, in a schematic representation of histology results of a titanium oxide substrate implanted subcutaneously. In the figure, "mg" means multinucleated giant cells and "ob" means osteoblasts. The substrate represented in Fig. 12B was implanted together with morphogen (OP-1) and newly formed osteoblasts are evident surrounding the substrate. By contrast, the substrate represented in Fig. 12A was implanted without morphogen and extensive multinucleated giant cell formation is evident surrounding the substrate. Accordingly, the morphogens' effect in inhibiting excessive bone mass loss in a mammal also may include inhibiting activation of these cells.

Example 13. Effect of Morphogen Neutralization on Bone Growth

[0114] The effect of the morphogens described herein on bone growth in developing mammals also may be evaluated using neutralizing antibodies specific for particular morphogens and assessing the effect of these antibodies on bone development. Specifically, anti-morphogen monoclonal and/or polyclonal antibodies may be prepared using standard methodologies including, for example, the protocol provided in Example 14, below.

[0115] Purified antibodies then are provided regularly to new born mice, e.g., 10-100µg/injection/day for 10-15 days. At 10 or 21 days, the mice are sacrificed and the effect of morphogen on bone development assessed by body weight, gross visual examination and histology. In this example, anti-OP-1 antibodies were used. Morphogen neutralization significantly stunted body growth, including bone growth, as indicated by the reduced body weight and reduced bone length of the treated mammals.

[0116] Similarly, morphogen activity may be assessed in fetal development in the mouse model using the following assay. Single lip injections comprising 10-100µg/injection of morphogen-specific antibody are administered to pregnant female mice during each day of the gestation period and bone development in treated and control new mice evaluated by standard histomorphometric analysis at birth. Similarly, single lip injections also may be provided to juvenile and adult mice (e.g., 10-100 µg) over a prolonged time (e.g., 10-15 days) to evaluate the effect on bone growth and bone integrity and to evaluate the onset of osteoporosis. The antibodies are anticipated to inhibit tissue morphogenesis, including bone growth and bone development, in the developing embryos.

Example 14. Screening Assay for Candidate Compounds which Alter Endogenous Morphogen Levels

[0117] Candidate compound(s) which may be administered to affect the level of a given morphogen may be found

using the following screening assay, in which the level of morphogen production by a cell type which produces measurable levels of the morphogen is determined with and without incubating the cell in culture with the compound, in order to assess the effects of the compound on the cell. This can be accomplished by detection of the morphogen either at the protein or RNA level.

14.1 Growth of Cells in Culture

[0118] Cell cultures of kidney, adrenals, urinary bladder, brain, or other organs, may be prepared as described widely in the literature. For example, kidneys may be explanted from neonatal or new born or young or adult rodents (mouse or rat) and used in organ culture as whole or sliced (1-4 mm) tissues. Primary tissue cultures and established cell lines, also derived from kidney, adrenals, urinary, bladder, brain, mammary, or other tissues may be established in multiwell plates (6 well or 24 well) according to conventional cell culture techniques, and are cultured in the absence or presence of serum for a period of time (1-7 days). Cells may be cultured, for example, in Dulbecco's Modified Eagle medium (Gibco, Long Island, NY) containing serum (e.g., fetal calf serum at 1%-10%, Gibco) or in serum-deprived medium, as desired, or in defined medium (e.g., containing insulin, transferrin, glucose, albumin, or other growth factors).

[0119] Samples for testing the level of morphogen production includes culture supernatants or cell lysates, collected periodically and evaluated for morphogen production by immunoblot analysis (Sambrook et al., eds., 1989, Molecular Cloning, Cold Spring Harbor Press, Cold Spring Harbor, NY), or a portion of the cell culture itself, collected periodically and used to prepare polyA+ RNA for RNA analysis. To monitor *de novo* morphogen synthesis, some cultures are labeled according to conventional procedures with an ^{35}S -methionine/ ^{35}S -cysteine mixture for 6-24 hours and then evaluated for morphogenic protein synthesis by conventional immunoprecipitation methods.

14.2 Determination of Level of Morphogenic Protein

[0120] In order to quantitate the production of a morphogenic protein by a cell type, an immunoassay may be performed to detect the morphogen using a polyclonal or monoclonal antibody specific for that protein. For example, OP-1 may be detected using a polyclonal antibody specific for OP-1 in an ELISA, as follows.

[0121] 1 μg /100 μl of affinity-purified polyclonal rabbit IgG specific for OP-1 is added to each well of a 96-well plate and incubated at 37°C for an hour. The wells are washed four times with 0.167M sodium borate buffer with 0.15 M NaCl (BSB), pH 8.2, containing 0.1% Tween 20. To minimize non-specific binding, the wells are blocked by filling completely with 1% bovine serum albumin (BSA) in BSB and incubating for 1 hour at 37°C. The wells are then washed four times with BSB containing 0.1% Tween 20. A 100 μl aliquot of an appropriate dilution of each of the test samples of cell culture supernatant is added to each well in triplicate and incubated at 37°C for 30 min. After incubation, 100 μl biotinylated rabbit anti-OP-1 serum (stock solution is about 1 mg/ml and diluted 1:400 in BSB containing 1% BSA before use) is added to each well and incubated at 37°C for 30 min. The wells are then washed four times with BSB containing 0.1% Tween 20. 100 μl streptavidin-alkaline (Southern Biotechnology Associates, Inc. Birmingham, Alabama, diluted 1:2000 in BSB containing 0.1% Tween 20 before use) is added to each well and incubated at 37°C for 30 min. The plates are washed four times with 0.5M Tris buffered Saline (TBS), pH 7.2. 50 μl substrate (ELISA Amplification System Kit, Life Technologies, Inc., Bethesda, MD) is added to each well incubated at room temperature for 15 min. Then, 50 μl amplifier (from the same amplification system kit) is added and incubated for another 15 min at room temperature. The reaction is stopped by the addition of 50 μl 0.3 M sulphuric acid. The OD at 490 nm of the solution in each well is recorded. To quantitate OP-1 in culture media, a OP-1 standard curve is performed in parallel with the test samples.

[0122] Polyclonal antibody may be prepared as follows. Each rabbit is given a primary immunization of 100 μg /500 μl *E. coli*-produced OP-1 monomer (amino acids 328-431 in SEQ ID NO:5) in 0.1% SDS mixed with 500 μl Complete Freund's Adjuvant. The antigen is injected subcutaneously at multiple sites on the back and flanks of the animal. The rabbit is boosted after a month in the same manner using incomplete Freund's Adjuvant. Test bleeds are taken from the ear vein seven days later. Two additional boosts and test bleeds are performed at monthly intervals until antibody against OP-1 is detected in the serum using an ELISA assay. Then, the rabbit is boosted monthly with 100 μg of antigen and bled (15 ml per bleed) at days seven and ten after boosting.

[0123] Monoclonal antibody specific for a given morphogen may be prepared as follows. A mouse is given two injections of *E. coli* produced OP-1 monomer. The first injection contains 100 μg of OP-1 in complete Freund's adjuvant and is given subcutaneously. The second injection contains 50 μg of OP-1 in incomplete adjuvant and is given intraperitoneally. The mouse then receives a total of 230 μg of OP-1 (amino acids 307-431 in SEQ ID NO:5) in four intraperitoneal injections at various times over an eight month period. One week prior to fusion, both mice are boosted intraperitoneally with 100 μg of OP-1 (307-431) and 30 μg of the N-terminal peptide (Ser₂₉₃-Asn₃₀₉-Cys) conjugated through the added cysteine to bovine serum albumin with SMCC crosslinking agent. This boost was repeated five days (IP), four days (IP), three days (IP) and one day (IV) prior to fusion. The mouse spleen cells are then fused to com-

mercially available myeloma cells at a ratio of 1:1 using PEG 1500 (Boeringer Mannheim, Germany), and the cell fusion is plated and screened for OP-1-specific antibodies using OP-1 (307-431) as antigen. The cell fusion and monoclonal screening then are according to standard procedures well described in standard texts widely available in the art.

5 SEQUENCE LISTING

[0124]

10 (1) GENERAL INFORMATION:

(i) APPLICANTS: Thangavel Kuberasampath
Charles Cohen
Hermann Oppermann
Engin Ozkayanak
15 David C. Rueger
Roy H.L. Pang

(ii) TITLE OF INVENTION: TREATMENT TO PREVENT LOSS OF AND/OR INCREASE BONE MASS IN
20 METABOLIC BONE DISEASE

(iii) NUMBER OF SEQUENCES: 33

(iv) CORRESPONDENCE ADDRESS:

25 (A) ADDRESSEE: Testa, Hurwitz & Thibault
(B) STREET: 53 State Street
(C) CITY: Boston
(D) STATE: Massachusetts
(E) COUNTRY: U.S.A.
30 (F) ZIP: 02109

(v) COMPUTER READABLE FORM:

35 (A) MEDIUM TYPE: Floppy Disk
(B) COMPUTER: IBM XT
(C) OPERATING SYSTEM: DOS 3.30
(D) SOFTWARE: PatentIn Release 1.0, Version 1.25

(vi) CURRENT APPLICATION DATA:

40 (B) FILING DATE:

(vii) PRIOR APPLICATION DATA:

45 (A) APPLICATION NUMBER: US 752,857
(B) FILING DATE: 30-AUG-1991

(viii) PRIOR APPLICATION DATA:

50 (A) APPLICATION NUMBER: US 667,274
(B) FILING DATE: 11-MAR-1991

(2) INFORMATION FOR SEQ ID NO:1:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 97 amino acids
(B) TYPE: amino acids

(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: Generic Sequence 1

(D) OTHER INFORMATION: Each Xaa indicates one of the 20 naturally-occurring L-isomer, α -amino acids or a derivative thereof.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

                                Xaa Xaa Xaa Xaa Xaa Xaa
                                1      5
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      10      15
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa
      20      25
Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      30      35
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      40      45      50
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
      55      60
Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      65      70

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      75      80
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
      85      90
Xaa Cys Xaa
      95

```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 97 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: Generic Sequence 2

(D) OTHER INFORMATION: Each Xaa indicates one of the 20 naturally-occurring L-isomer, α -amino acids or a derivative thereof.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

5

10

15

20

25

30

35

40

```

                Xaa Xaa Xaa Xaa Xaa Xaa
                  1             5
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
          10             15
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa
          20             25
Cys Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa
          30             35

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
40             45             50
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
          55             60
Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
          65             70
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
          75             80
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
          85             90
Xaa Cys Xaa
          95

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 97 amino acids
- (B) TYPE: amino acids
- (C) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

55

- (A) NAME: Generic Sequence 3
- (D) OTHER INFORMATION: wherein each Xaa is independently selected from a group of one or more specified amino acids as defined in the specification.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Leu Tyr Val Xaa Phe
1 5

5

10

Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
10

Xaa Ala Pro Gly Xaa Xaa Xaa Ala
15 20

15

Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
25 30

Xaa Pro Xaa Xaa Xaa Xaa Xaa
35

20

Xaa Xaa Xaa Asn His Ala Xaa Xaa
40 45

Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa
50

25

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
55 60

30

Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
65

Xaa Xaa Xaa Leu Xaa Xaa Xaa
70 75

35

Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
80

Xaa Xaa Xaa Xaa Met Xaa Val Xaa
85 90

40

Xaa Cys Gly Cys Xaa
95

45

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

50

(A) LENGTH: 102 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

55

(ix) FEATURE:

(A) NAME: Generic Sequence 4

(D) OTHER INFORMATION: wherein each Xaa is independently selected from a group of one or more specified amino acids as defined in the specification.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

5

10

15

20

25

30

35

40

45

```

Cys Xaa Xaa Xaa Xaa Leu Tyr Val Xaa Phe
  1               5               10
Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
              15
Xaa Ala Pro Xaa Gly Xaa Xaa Ala
  20               25
Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
              30               35
Xaa Pro Xaa Xaa Xaa Xaa Xaa
              40
Asn Xaa Xaa Asn His Ala Xaa Xaa
              45               50
Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa
              55
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
              60               65
Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
              70
Xaa Xaa Xaa Leu Xaa Xaa Xaa
              75               80
Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
              85
Xaa Xaa Xaa Xaa Met Xaa Val Xaa
              90               95
Xaa Cys Gly Cys Xaa
              100

```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

50

(A) LENGTH: 139 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: hOP-1 (mature form)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

5		Ser	Thr	Gly	Ser	Lys	Gln	Arg	Ser	Gln
	1					5				
		Asn	Arg	Ser	Lys	Thr	Pro	Lys	Asn	Gln
10	10					15				
		Glu	Ala	Leu	Arg	Met	Ala	Asn	Val	Ala
	20					25				
15		Glu	Asn	Ser	Ser	Ser	Asp	Gln	Arg	Gln
	30					35				
		Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val
	40					45				
20		Ser	Phe	Arg	Asp	Leu	Gly	Trp	Gln	Asp
	50					55				
		Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala
25	55					60				
		Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala
	65					70				
		Phe	Pro	Leu	Asn	Ser	Tyr	Met	Asn	Ala
30	75					80				
		Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu
	85					90				
35										
		Val	His	Phe	Ile	Asn	Pro	Glu	Thr	Val
	95									
40		Pro	Lys	Pro	Cys	Cys	Ala	Pro	Thr	Gln
	100					105				
		Leu	Asn	Ala	Ile	Ser	Val	Leu	Tyr	Phe
45	110					115				
		Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	Lys
	120					125				
		Lys	Tyr	Arg	Asn	Met	Val	Val	Arg	Ala
50	130					135				
		Cys	Gly	Cys	His					

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 139 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: mOP-1 (mature form)

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

15

Ser	Thr	Gly	Gly	Lys	Gln	Arg	Ser	Gln
1				5				

Asn	Arg	Ser	Lys	Thr	Pro	Lys	Asn	Gln
10					15			

20

Glu	Ala	Leu	Arg	Met	Ala	Ser	Val	Ala
	20					25		

Glu	Asn	Ser	Ser	Ser	Asp	Gln	Arg	Gln
		30					35	

25

30

35

40

45

50

55

	Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val
				40					45
5	Ser	Phe	Arg	Asp	Leu	Gly	Trp	Gln	Asp
				50					
	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala
10	55					60			
	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala
	65						70		
	Phe	Pro	Leu	Asn	Ser	Tyr	Met	Asn	Ala
15			75					80	
	Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu
			85						90
20	Val	His	Phe	Ile	Asn	Pro	Asp	Thr	Val
				95					
	Pro	Lys	Pro	Cys	Cys	Ala	Pro	Thr	Gln
25	100					105			
	Leu	Asn	Ala	Ile	Ser	Val	Leu	Tyr	Phe
	110						115		
	Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	Lys
30			120					125	
	Lys	Tyr	Arg	Asn	Met	Val	Val	Arg	Ala
			130						135
35	Cys	Gly	Cys	His					

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 139 amino acids
 (B) TYPE: amino acids
 (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME: hOP-2 (mature form)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

	Ala	Val	Arg	Pro	Leu	Arg	Arg	Arg	Gln
	1				5				
5	Pro	Lys	Lys	Ser	Asn	Glu	Leu	Pro	Gln
	10					15			
	Ala	Asn	Arg	Leu	Pro	Gly	Ile	Phe	Asp
		20					25		
10	Asp	Val	His	Gly	Ser	His	Gly	Arg	Gln
			30					35	
	Val	Cys	Arg	Arg	His	Glu	Leu	Tyr	Val
15				40					45
	Ser	Phe	Gln	Asp	Leu	Gly	Trp	Leu	Asp
					50				
20	Trp	Val	Ile	Ala	Pro	Gln	Gly	Tyr	Ser
	55					60			
	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ser
		65					70		
25	Phe	Pro	Leu	Asp	Ser	Cys	Met	Asn	Ala
			75					80	
	Thr	Asn	His	Ala	Ile	Leu	Gln	Ser	Leu
30				85					90
	Val	His	Leu	Met	Lys	Pro	Asn	Ala	Val
					95				
35	Pro	Lys	Ala	Cys	Cys	Ala	Pro	Thr	Lys
	100					105			
	Leu	Ser	Ala	Thr	Ser	Val	Leu	Tyr	Tyr
		110					115		
40	Asp	Ser	Ser	Asn	Asn	Val	Ile	Leu	Arg
			120					125	
	Lys	His	Arg	Asn	Met	Val	Val	Lys	Ala
				130					135
45	Cys	Gly	Cys	His					

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 139 amino acids
 (B) TYPE: amino acids
 (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: mOP-2 (mature form)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

5		Ala	Ala	Arg	Pro	Leu	Lys	Arg	Arg	Gln
		1				5				
10		Pro	Lys	Lys	Thr	Asn	Glu	Leu	Pro	His
		10					15			
		Pro	Asn	Lys	Leu	Pro	Gly	Ile	Phe	Asp
			20					25		
15		Asp	Gly	His	Gly	Ser	Arg	Gly	Arg	Glu
				30					35	
		Val	Cys	Arg	Arg	His	Glu	Leu	Tyr	Val
20					40					45
		Ser	Phe	Arg	Asp	Leu	Gly	Trp	Leu	Asp
						50				
		Trp	Val	Ile	Ala	Pro	Gln	Gly	Tyr	Ser
25							60			
		Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala
			65					70		
30		Phe	Pro	Leu	Asp	Ser	Cys	Met	Asn	Ala
				75					80	
		Thr	Asn	His	Ala	Ile	Leu	Gln	Ser	Leu
35					85					90
		Val	His	Leu	Met	Lys	Pro	Asp	Val	Val
40						95				
		Pro	Lys	Ala	Cys	Cys	Ala	Pro	Thr	Lys
		100					105			
		Leu	Ser	Ala	Thr	Ser	Val	Leu	Tyr	Tyr
45			110					115		
		Asp	Ser	Ser	Asn	Asn	Val	Ile	Leu	Arg
				120					125	
50		Lys	His	Arg	Asn	Met	Val	Val	Lys	Ala
					130					135
		Cys	Gly	Cys	His					

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 96 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: CBMP2A(fx)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Cys Lys Arg His Pro Leu Tyr Val Asp Phe Ser
  1             5             10
Asp Val Gly Trp Asn Asp Trp Ile Val Ala Pro
      15             20
Pro Gly Tyr His Ala Phe Tyr Cys His Gly Glu
      25             30
Cys Pro Phe Pro Leu Ala Asp His Leu Asn Ser
      35             40

Thr Asn His Ala Ile Val Gln Thr Leu Val Asn
  45             50             55
Ser Val Asn Ser Lys Ile Pro Lys Ala Cys Cys
      60             65
Val Pro Thr Glu Leu Ser Ala Ile Ser Met Leu
      70             75
Tyr Leu Asp Glu Asn Glu Lys Val Val Leu Lys
      80             85
Asn Tyr Gln Asp Met Val Val Glu Gly Cys Gly
      90             95
Cys Arg
100

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 101 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: CBMP2B(fx)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

5 Cys Arg Arg His Ser
 1 5
 Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asn
 10 10 15
 Asp Trp Ile Val Ala Pro Pro Gly Tyr Gln Ala
 20 25
 15 Phe Tyr Cys His Gly Asp Cys Pro Phe Pro Leu
 30 35
 20
 Ala Asp His Leu Asn Ser Thr Asn His Ala Ile
 40 45
 Val Gln Thr Leu Val Asn Ser Val Asn Ser Ser
 25 50 55 60
 Ile Pro Lys Ala Cys Cys Val Pro Thr Glu Leu
 65 70
 30 Ser Ala Ile Ser Met Leu Tyr Leu Asp Glu Tyr
 75 80
 Asp Lys Val Val Leu Lys Asn Tyr Gln Glu Met
 85 90
 35 Val Val Glu Gly Cys Gly Cys Arg
 95 100

40 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 102 amino acids
 (B) TYPE: amino acids
 (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

50 (ix) FEATURE:

(A) NAME: DPP(fx)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

55

Cys Arg Arg His Ser Leu Tyr Val Asp Phe Ser
 1 5 10
 5 Asp Val Gly Trp Asp Asp Trp Ile Val Ala Pro
 15 20
 Leu Gly Tyr Asp Ala Tyr Tyr Cys His Gly Lys
 25 30
 10
 15 Cys Pro Phe Pro Leu Ala Asp His Phe Asn Ser
 35 40
 Thr Asn His Ala Val Val Gln Thr Leu Val Asn
 45 50 55
 20 Asn Asn Asn Pro Gly Lys Val Pro Lys Ala Cys
 60 65
 Cys Val Pro Thr Gln Leu Asp Ser Val Ala Met
 70 75
 25 Leu Tyr Leu Asn Asp Gln Ser Thr Val Val Leu
 80 85
 30 Lys Asn Tyr Gln Glu Met Thr Val Val Gly Cys
 90 95
 Gly Cys Arg
 100
 35

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 102 amino acids
 (B) TYPE: amino acids
 (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME: Vgl(fx)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Cys Lys Lys Arg His Leu Tyr Val Glu Phe Lys
 1 5 10
 5 Asp Val Gly Trp Gln Asn Trp Val Ile Ala Pro
 15 20
 Gln Gly Tyr Met Ala Asn Tyr Cys Tyr Gly Glu
 25 30
 10
 Cys Pro Tyr Pro Leu Thr Glu Ile Leu Asn Gly
 35 40
 Ser Asn His Ala Ile Leu Gln Thr Leu Val His
 45 50 55
 20 Ser Ile Glu Pro Glu Asp Ile Pro Leu Pro Cys
 60 65
 Cys Val Pro Thr Lys Met Ser Pro Ile Ser Met
 70 75
 25 Leu Phe Tyr Asp Asn Asn Asp Asn Val Val Leu
 80 85
 Arg His Tyr Glu Asn Met Ala Val Asp Glu Cys
 90 95
 30 Gly Cys Arg
 100

35 (2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 102 amino acids
 (B) TYPE: amino acids
 (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45 (ix) FEATURE:

(A) NAME: Vgr-1(fx)

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Cys Lys Lys His Glu Leu Tyr Val Ser Phe Gln
 1 5 10
 5 Asp Val Gly Trp Gln Asp Trp Ile Ile Ala Pro
 15 20
 Xaa Gly Tyr Ala Ala Asn Tyr Cys Asp Gly Glu
 10 25 30

 15 Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala
 35 40
 Thr Asn His Ala Ile Val Gln Thr Leu Val His
 45 50 55
 20 Val Met Asn Pro Glu Tyr Val Pro Lys Pro Cys
 60 65
 Cys Ala Pro Thr Lys Val Asn Ala Ile Ser Val
 25 70 75
 Leu Tyr Phe Asp Asp Asn Ser Asn Val Ile Leu
 80 85
 30 Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys
 90 95
 Gly Cys His
 100
 35

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 106 amino acids
 (B) TYPE: protein
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- 50 (A) ORGANISM: human
 (F) TISSUE TYPE: BRAIN

(ix) FEATURE:

- 55 (D) OTHER INFORMATION: /product= "GDF-1 (fx)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Cys Arg Ala Arg Arg Leu Tyr Val Ser Phe Arg Glu Val Gly
 1 5 10
 5 Trp His Arg Trp Val Ile Ala Pro Arg Gly Phe Leu Ala Asn Tyr
 15 20 25
 Cys Gln Gly Gln Cys Ala Leu Pro Val Ala Leu Ser Gly Ser Gly
 30 35 40
 10
 Gly Pro Pro Ala Leu Asn His Ala Val Leu Arg Ala Leu Met His
 45 50 55
 15 Ala Ala Ala Pro Gly Ala Ala Asp Leu Pro Cys Cys Val Pro Ala
 60 65 70
 20 Arg Leu Ser Pro Ile Ser Val Leu Phe Phe Asp Asn Ser Asp Asn
 75 80 85
 Val Val Leu Arg Gln Tyr Glu Asp Met Val Val Asp Glu Cys Gly
 90 95 100
 25 Cys Arg
 105

(2) INFORMATION FOR SEQ ID NO:15:

30 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

40

Cys Xaa Xaa Xaa Xaa
 1 5

45 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1822 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

55

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HOMO SAPIENS

(F) TISSUE TYPE: HIPPOCAMPUS

(ix) FEATURE:

- 5 (A) NAME/KEY: CDS
 (B) LOCATION: 49..1341
 (D) OTHER INFORMATION:/standard_name= "hOP1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

10

GGTGCGGGCC CGGAGCCCGG AGCCCGGGTA GCGCGTAGAG CCGGCGCG ATG CAC GTG 57
 Met His Val
 1

15

20

25

30

35

40

45

50

55

	CGC	TCA	CTG	CGA	GCT	GCG	GCG	CCG	CAC	AGC	TTC	GTG	GCG	CTC	TGG	GCA	105
	Arg	Ser	Leu	Arg	Ala	Ala	Ala	Pro	His	Ser	Phe	Val	Ala	Leu	Trp	Ala	
		5					10					15					
5	CCC	CTG	TTC	CTG	CTG	GCG	TCC	GCC	CTG	GCC	GAC	TTC	AGC	CTG	GAC	AAC	153
	Pro	Leu	Phe	Leu	Leu	Arg	Ser	Ala	Leu	Ala	Asp	Phe	Ser	Leu	Asp	Asn	
	20					25					30					35	
10	GAG	GTG	CAC	TCG	AGC	TTC	ATC	CAC	CGG	CGC	CTC	CGC	AGC	CAG	GAG	CGG	201
	Glu	Val	His	Ser	Ser	Phe	Ile	His	Arg	Arg	Leu	Arg	Ser	Gln	Glu	Arg	
					40					45					50		
15	CGG	GAG	ATG	CAG	CGC	GAG	ATC	CTC	TCC	ATT	TTG	GGC	TTG	CCC	CAC	CGC	249
	Arg	Glu	Met	Gln	Arg	Glu	Ile	Leu	Ser	Ile	Leu	Gly	Leu	Pro	His	Arg	
				55					60					65			
20	CCG	CGC	CCG	CAC	CTC	CAG	GGC	AAG	CAC	AAC	TCG	GCA	CCC	ATG	TTC	ATG	297
	PrBo	Arg	Pro	His	Leu	Gln	Gly	Lys	His	Asn	Ser	Ala	Pro	Met	Phe	Met	
				70				75					80				
25	CTG	GAC	CTG	TAC	AAC	GCC	ATG	GCG	GTG	GAG	GAG	GGC	GGC	GGG	CCC	GGC	345
	Leu	Asp	Leu	Tyr	Asn	Ala	Met	Ala	Val	Glu	Glu	Gly	Gly	Gly	Pro	Gly	
		85					90					95					
30	GGC	CAG	GGC	TTC	TCC	TAC	CCC	TAC	AAG	GCC	GTC	TTC	AGT	ACC	CAG	GGC	393
	Gly	Gln	Gly	Phe	Ser	Tyr	Pro	Tyr	Lys	Ala	Val	Phe	Ser	Thr	Gln	Gly	
	100					105					110					115	
35	CCC	CCT	CTG	GCC	AGC	CTG	CAA	GAT	AGC	CAT	TTC	CTC	ACC	GAC	GCC	GAC	441
	Pro	Pro	Leu	Ala	Ser	Leu	Gln	Asp	Ser	His	Phe	Leu	Thr	Asp	Ala	Asp	
					120					125					130		
40	ATG	GTC	ATG	AGC	TTC	GTC	AAC	CTC	GTG	GAA	CAT	GAC	AAG	GAA	TTC	TTC	489
	Met	Val	Met	Ser	Phe	Val	Asn	Leu	Val	Glu	His	Asp	Lys	Glu	Phe	Phe	
				135				140						145			
45	CAC	CCA	CGC	TAC	CAC	CAT	CGA	GAG	TTC	CGG	TTT	GAT	CTT	TCC	AAG	ATC	537
	His	Pro	Arg	Tyr	His	His	Arg	Glu	Phe	Arg	Phe	Asp	Leu	Ser	Lys	Ile	
				150				155					160				
50	CCA	GAA	GGG	GAA	GCT	GTC	ACG	GCA	GCC	GAA	TTC	CGG	ATC	TAC	AAG	GAC	585
	Pro	Glu	Gly	Glu	Ala	Val	Thr	Ala	Ala	Glu	Phe	Arg	Ile	Tyr	Lys	Asp	
		165					170					175					
55	TAC	ATC	CGG	GAA	CGC	TTC	GAC	AAT	GAG	ACG	TTC	CGG	ATC	AGC	GTT	TAT	633
	Tyr	Ile	Arg	Glu	Arg	Phe	Asp	Asn	Glu	Thr	Phe	Arg	Ile	Ser	Val	Tyr	
	180					185					190					195	
60	CAG	GTG	CTC	CAG	GAG	CAC	TTG	GGC	AGG	GAA	TCG	GAT	CTC	TTC	CTG	CTC	681
	Gln	Val	Leu	Gln	Glu	His	Leu	Gly	Arg	Glu	Ser	Asp	Leu	Phe	Leu	Leu	
					200					205					210		

GAC Asp	AGC Ser	CGT Arg	ACC Thr 215	CTC Leu	TGG Trp	GCC Ala	TCG Ser	GAG Glu	GAG Glu	GGC Gly	TGG Trp	CTG Leu	GTG Val	TTT Phe	GAC Asp	729
ATC Ile	ACA Thr	GCC Ala 230	ACC Thr	AGC Ser	AAC Asn	CAC His	TGG Trp 235	GTG Val	GTC Val	AAT Asn	CCG Pro	CGG Arg 240	CAC His	AAC Asn	CTG Leu	777
GGC Gly	CTG Leu 245	CAG Gln	CTC Leu	TCG Ser	GTG Val	GAG Glu 250	ACG Thr	CTG Leu	GAT Asp	GGG Gly	CAG Gln 255	AGC Ser	ATC Ile	AAC Asn	CCC Pro	825
AAG Lys 260	TTG Leu	GCG Ala	GGC Gly	CTG Leu	ATT Ile 265	GGG Gly	CGG Arg	CAC His	GGG Gly	CCC Pro 270	CAG Gln	AAC Asn	AAG Lys	CAG Gln	CCC Pro 275	873
TTC Phe	ATG Met	GTG Val	GCT Ala	TTC Phe 280	TTC Phe	AAG Lys	GCC Ala	ACG Thr	GAG Glu 285	GTC Val	CAC His	TTC Phe	CGC Arg	AGC Ser 290	ATC Ile	921
CGG Arg	TCC Ser	ACG Thr	GGG Gly 295	AGC Ser	AAA Lys	CAG Gln	CGC Arg	AGC Ser 300	CAG Gln	AAC Asn	CGC Arg	TCC Ser	AAG Lys 305	ACG Thr	CCC Pro	969
AAG Lys	AAC Asn	CAG Gln 310	GAA Glu	GCC Ala	CTG Leu	CGG Arg	ATG Met 315	GCC Ala	AAC Asn	GTG Val	GCA Ala 320	GAG Glu	AAC Asn	AGC Ser	AGC Ser	1017
AGC Ser	GAC Asp 325	CAG Gln	AGG Arg	CAG Gln	GCC Ala	TGT Cys 330	AAG Lys	AAG Lys	CAC His	GAG Glu	CTG Leu 335	TAT Tyr	GTC Val	AGC Ser	TTC Phe	1065
CGA Arg 340	GAC Asp	CTG Leu	GGC Gly	TGG Trp	CAG Gln 345	GAC Asp	TGG Trp	ATC Ile	ATC Ile	GCG Ala 350	CCT Pro	GAA Glu	GGC Gly	TAC Tyr	GCC Ala 355	1113
GCC Ala	TAC Tyr	TAC Tyr	TGT Cys 360	GAG Glu	GGG Gly	GAG Glu	TGT Cys	GCC Ala 365	TTC Phe	CCT Pro	CTG Leu	AAC Asn	TCC Ser	TAC Tyr 370	ATG Met	1161
AAC Asn	GCC Ala	ACC Thr 375	AAC Asn	CAC His	GCC Ala	ATC Ile	GTG Val	CAG Gln 380	ACG Thr	CTG Leu	GTC Val	CAC His	TTC Phe 385	ATC Ile	AAC Asn	1209
CCG Pro	GAA Glu	ACG Thr 390	GTG Val	CCC Pro	AAG Lys	CCC Pro	TGC Cys 395	TGT Cys	GCG Ala	CCC Pro	ACG Thr 400	CAG Gln	CTC Leu	AAT Asn	GCC Ala	1257
ATC Ile	TCC Ser 405	GTC Val	CTC Leu	TAC Tyr	TTC Phe	GAT Asp 410	GAC Asp	AGC Ser	TCC Ser	AAC Asn	GTC Val 415	ATC Ile	CTG Leu	AAG Lys	AAA Lys	1305

TAC AGA AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCCTCC 1351
 Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430

5

GAGAATTCAG ACCCTTTGGG GCCAAGTTTT TCTGGATCCT CCATTGCTCG CCTTGGCCAG 1411
 GAACCAGCAG ACCAACTGCC TTTTGTGAGA CCTTCCCCTC CCTATCCCCA ACTTTAAAGG 1471
 10 TGTGAGAGTA TTAGGAAACA TGAGCAGCAT ATGGCTTTTG ATCAGTTTTT CAGTGGCAGC 1531
 ATCCAATGAA CAAGATCCTA CAAGCTGTGC AGGCAAAACC TAGCAGGAAA AAAAAACAAC 1591
 GCATAAAGAA AAATGGCCGG GCCAGGTCAT TGGCTGGGAA GTCTCAGCCA TGCACGGACT 1651
 15 CGTTTCCAGA GGTAATTATG AGCGCCTACC AGCCAGGCCA CCCAGCCGTG GGAGGAAGGG 1711
 GGCGTGGCAA GGGGTGGGCA CATTGGTGTC TGTGCGAAAG GAAAATTGAC CCGGAAGTTC 1771
 CTGTAATAAA TGTCACAATA AAACGAATGA ATGAAAAAAA AAAAAAAAAA A 1822
 20

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 431 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(D) OTHER INFORMATION: /Product="OP1-PP"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
 1 5 10 15
 Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
 20 25 30
 Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
 35 40 45
 Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
 50 55 60
 Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
 65 70 75 80
 Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly
 85 90 95

Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser
 100 105 110
 5 Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr
 115 120 125
 Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys
 130 135 140
 10 Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu
 145 150 155 160
 Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile
 15 165 170 175
 Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile
 180 185 190
 20 Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu
 195 200 205
 Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu
 210 215 220
 25 Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg
 225 230 235 240
 His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser
 245 250 255
 30 Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn
 260 265 270
 Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe
 275 280 285
 35 Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser
 290 295 300
 40 Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu
 305 310 315 320
 Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr
 325 330 335
 45 V|Bal Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu
 340 345 350
 Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn
 355 360 365
 50
 55

Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His
 370 375 380

5 Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln
 385 390 395 400

Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile
 405 410 415

10 Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1873 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HURIDAE
 (F) TISSUE TYPE: EMBRYO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 104..1393
 (D) OTHER INFORMATION: /note= "MOP1 (CDNA)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTGCAGCAAG TGACCTCGGG TCGTGGACCG CTGCCCTGCC CCCTCCGCTG CCACCTGGGG 60

CGGCGCGGGC CCGGTGCCCC GGATCGCGCG TAGAGCCGGC GCG ATG CAC GTG CGC 115
 Met His Val Arg
 1

45 TCG CTG CGC GCT GCG GCG CCA CAC AGC TTC GTG GCG CTC TGG GCG CCT 163
 Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala Pro
 5 10 15 20

CTG TTC TTG CTG CGC TCC GCC CTG GCC GAT TTC AGC CTG GAC AAC GAG 211
 Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser Leu Asp Asn Glu
 25 30 35

GTG CAC TCC AGC TTC ATC CAC CGG CGC CTC CGC AGC CAG GAG CGG CGG 259
 Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser Gln Glu Arg Arg
 40 45 50

	GAG	ATG	CAG	CGG	GAG	ATC	CTG	TCC	ATC	TTA	GGG	TTG	CCC	CAT	CGC	CCG	307
	Glu	Met	Gln	Arg	Glu	Ile	Leu	Ser	Ile	Leu	Gly	Leu	Pro	His	Arg	Pro	
			55				60					65					
5																	
	CGC	CCG	CAC	CTC	CAG	GGA	AAG	CAT	AAT	TCG	GCG	CCC	ATG	TTC	ATG	TTG	355
	Arg	Pro	His	Leu	Gln	Gly	Lys	His	Asn	Ser	Ala	Pro	Met	Phe	Met	Leu	
		70					75					80					
10																	
	GAC	CTG	TAC	AAC	GCC	ATG	GCG	GTG	GAG	GAG	AGC	GGG	CCG	GAC	GGA	CAG	403
	Asp	Leu	Tyr	Asn	Ala	Met	Ala	Val	Glu	Glu	Ser	Gly	Pro	Asp	Gly	Gln	
	85					90					95					100	
15																	
	GGC	TTC	TCC	TAC	CCC	TAC	AAG	GCC	GTC	TTC	AGT	ACC	CAG	GGC	CCC	CCT	451
	Gly	Phe	Ser	Tyr	Pro	Tyr	Lys	Ala	Val	Phe	Ser	Thr	Gln	Gly	Pro	Pro	
					105					110					115		
20																	
	TTA	GCC	AGC	CTG	CAG	GAC	AGC	CAT	TTC	CTC	ACT	GAC	GCC	GAC	ATG	GTC	499
	Leu	Ala	Ser	Leu	Gln	Asp	Ser	His	Phe	Leu	Thr	Asp	Ala	Asp	Met	Val	
					120					125				130			
25																	
	ATG	AGC	TTC	GTC	AAC	CTA	GTG	GAA	CAT	GAC	AAA	GAA	TTC	TTC	CAC	CCT	547
	Met	Ser	Phe	Val	Asn	Leu	Val	Glu	His	Asp	Lys	Glu	Phe	Phe	His	Pro	
			135					140					145				
	CGA	TAC	CAC	CAT	CGG	GAG	TTC	CGG	TTT	GAT	CTT	TCC	AAG	ATC	CCC	GAG	595
	Arg	Tyr	His	His	Arg	Glu	Phe	Arg	Phe	Asp	Leu	Ser	Lys	Ile	Pro	Glu	
		150					155					160					
30																	
	GGC	GAA	CGG	GTG	ACC	GCA	GCC	GAA	TTC	AGG	ATC	TAT	AAG	GAC	TAC	ATC	643
	Gly	Glu	Arg	Val	Thr	Ala	Ala	Glu	Phe	Arg	Ile	Tyr	Lys	Asp	Tyr	Ile	
	165					170					175					180	
35																	
	CGG	GAG	CGA	TTT	GAC	AAC	GAG	ACC	TTC	CAG	ATC	ACA	GTC	TAT	CAG	GTG	691
	Arg	Glu	Arg	Phe	Asp	Asn	Glu	Thr	Phe	Gln	Ile	Thr	Val	Tyr	Gln	Val	
					185					190					195		
40																	
	CTC	CAG	GAG	CAC	TCA	GGC	AGG	GAG	TCG	GAC	CTC	TTC	TTG	CTG	GAC	AGC	739
	Leu	Gln	Glu	His	Ser	Gly	Arg	Glu	Ser	Asp	Leu	Phe	Leu	Leu	Asp	Ser	
				200					205					210			
45																	
	CGC	ACC	ATC	TGG	GCT	TCT	GAG	GAG	GGC	TGG	TTG	GTG	TTT	GAT	ATC	ACA	787
	Arg	Thr	Ile	Trp	Ala	Ser	Glu	Glu	Gly	Trp	Leu	Val	Phe	Asp	Ile	Thr	
			215					220					225				
50																	
	GCC	ACC	AGC	AAC	CAC	TGG	GTG	GTC	AAC	CCT	CGG	CAC	AAC	CTG	GGC	TTA	835
	Ala	Thr	Ser	Asn	His	Trp	Val	Val	Asn	Pro	Arg	His	Asn	Leu	Gly	Leu	
		230					235					240					
55																	
	CAG	CTC	TCT	GTG	GAG	ACC	CTG	GAT	GGG	CAG	AGC	ATC	AAC	CCC	AAG	TTG	883
	Gln	Leu	Ser	Val	Glu	Thr	Leu	Asp	Gly	Gln	Ser	Ile	Asn	Pro	Lys	Leu	
	245					250					255					260	

	GCA GGC CTG ATT GGA CGG CAT GGA CCC CAG AAC AAG CAA CCC TTC ATG Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn Lys Gln Pro Phe Met 265 270 275	931
5	GTG GCC TTC TTC AAG GCC ACG GAA GTC CAT CTC CGT AGT ATC CGG TCC Val Ala Phe Phe Lys Ala Thr Glu Val His Leu Arg Ser Ile Arg Ser 280 285 290	979
10	ACG GGG GGC AAG CAG CGC AGC CAG AAT CGC TCC AAG ACG CCA AAG AAC Thr Gly Gly Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro Lys Asn 295 300 305	1027
15	CAA GAG GCC CTG AGG ATG GCC AGT GTG GCA GAA AAC AGC AGC AGT GAC Gln Glu Ala Leu Arg Met Ala Ser Val Ala Glu Asn Ser Ser Ser Asp 310 315 320	1075
20	CAG AGG CAG GCC TGC AAG AAA CAT GAG CTG TAC GTC AGC TTC CGA GAC Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg Asp 325 330 335 340	1123
25	CTT GGC TGG CAG GAC TGG ATC ATT GCA CCT GAA GGC TAT GCT GCC TAC Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala Tyr 345 350 355	1171
30	TAC TGT GAG GGA GAG TGC GCC TTC CCT CTG AAC TCC TAC ATG AAC GCC Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn Ala 360 365 370	1219
35	ACCB AAC CAC GCC ATC GTC CAG ACA CTG GTT CAC TTC ATC AAC CCA GAC Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn Pro Asp 375 380 385	1267
40	ACA GTA CCC AAG CCC TGC TGT GCG CCC ACC CAG CTC AAC GCC ATC TCT Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile Ser 390 395 400	1315
45	GTC CTC TAC TTC GAC GAC AGC TCT AAT GTC ATC CTG AAG AAG TAC AGA Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr Arg 405 410 415 420	1363
50	AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCTTCC TGAGACCCTG Asn Met Val Val Arg Ala Cys Gly Cys His 425 430	1413
55	ACCTTTGCGG GGCCACACCT TTCCAAATCT TCGATGTCTC ACCATCTAAG TCTCTCACTG	1473
	CCCACCTTGG CGAGGAGAAC AGACCAACCT CTCCTGAGCC TTCCCTCACC TCCCAACCGG	1533
	AAGCATGTAA GGGTTCCAGA AACCTGAGCG TGCAGCAGCT GATGAGCGCC CTTTCCTTCT	1593
	GGCACGTGAC GGACAAGATC CTACCAGCTA CCACAGCAAA CGCCTAAGAG CAGGAAAAAT	1653

GTCTGCCAGG AAAGTGTCCA GTGTCCACAT GGCCCTGGC GCTCTGAGTC TTTGAGGAGT 1713
 AATCGCAAGC CTCGTTTACG TGCAGCAGAA GGAAGGGCTT AGCCAGGGTG GGCGCTGGCG 1773
 TCTGTGTTGA AGGGAAACCA AGCAGAAGCC ACTGTAATGA TATGTCACAA TAAAACCCAT 1833
 GAATGAAAAA AAAAAAAAAA AAAAAAAAAA AAAAGAATTC 1873

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 430 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(D) OTHER INFORMATION: /product= "mOP1-PP"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
 1 5 10 15
 Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
 20 25 30
 Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
 35 40 45
 Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
 50 55 60
 Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
 65 70 75 80
 Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Ser Gly
 85 90 95
 Pro Asp Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr
 100 105 110
 Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp
 115 120 125
 Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu
 130 135 140
 Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu Ser
 145 150 155 160

Lys Ile Pro Glu Gly Glu Arg Val Thr Ala Ala Glu Phe Arg Ile Tyr
 165 170 175
 5 Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Gln Ile Thr
 180 185 190
 Val Tyr Gln Val Leu Gln Glu His Ser Gly Arg Glu Ser Asp Leu Phe
 195 200 205
 10 Leu Leu Asp Ser Arg Thr Ile Trp Ala Ser Glu Glu Gly Trp Leu Val
 210 215 220
 Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg His
 225 230 235 240
 15 Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser Ile
 245 250 255
 Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn Lys
 260 265 270
 Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Leu Arg
 275 280 285
 25 Ser Ile Arg Ser Thr Gly Gly Lys Gln Arg Ser Gln Asn Arg Ser Lys
 290 295 300
 Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Ser Val Ala Glu Asn
 305 310 315 320
 30 Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val
 325 330 335
 Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly
 340 345 350
 35 Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser
 355 360 365
 Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe
 370 375 380
 Ile Asn Pro Asp Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu
 385 390 395 400
 45 Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu
 405 410 415
 Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430
 50

(2) INFORMATION FOR SEQ ID NO:20:

(i)SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1723 base pairs
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii)MOLECULE TYPE: cDNA

(vi)ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(F) TISSUE TYPE: HIPPOCAMPUS

(ix)FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 490..1696

(D) OTHER INFORMATION: /note= "hOP2 (cDNA)"

(xi)SEQUENCE DESCRIPTION: SEQ ID NO:20:

20 GCGCCCGGCA GAGCAGGAGT GGCTGGAGGA GCTGTGGTTG GAGCAGGAGG TGGCACGGCA 60
 GGGCTGGAGG GCTCCCTATG AGTGGCGGAG ACGGCCCAGG AGGCGCTGGA GCAACAGCTC 120
 CCACACCGCA CCAAGCGGTG GCTGCAGGAG CTCGCCCATC GCCCCTGCGC TGCTCGGACC 180
 25 GCGGCCACAG CCGGACTGGC GGGTACGGCG GCGACAGAGG CATTGGCCGA GAGTCCCAGT 240
 CCGCAGAGTA GCCCCGGCCT CGAGGCGGTG GCGTCCCGGT CCTCTCCGTC CAGGAGCCAG 300
 GACAGGTGTC GCGCGGCGGG GCTCCAGGGA CCGCGCCTGA GGCCGGCTGC CCGCCCGTCC 360
 CGCCCCGCCC CGCCGCCCCG CGCCCGCCGA GCCCAGCCTC CTTGCCGTCC GGGCGTCCCC 420
 AGGCCCTGGG TCGGCCCGGG AGCCGATGCG CGCCCGCTGA GCGCCCCAGC TGAGCGCCCC 480
 35 CGGCCTGCC ATG ACC GCG CTC CCC GGC CCG CTC TGG CTC CTG GGC CTG 528
 Met Thr Ala Leu Pro Gly Pro Leu Trp Leu Leu Gly Leu
 1 5 10
 GCG CTA TGC GCG CTG GGC GGG GGC GGC CCC GGC CTG CGA CCC CCG CCC 576
 40 Ala Leu Cys Ala Leu Gly Gly Gly Gly Pro Gly Leu Arg Pro Pro Pro
 15 20 25
 GGC TGT CCC CAG CGA CGT CTG GGC GCG CGC GAG CGC CGG GAC GTG CAG 624
 45 Gly Cys Pro Gln Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Val Gln
 30 35 40 45
 CGC GAG ATC CTG GCG GTG CTC GGG CTG CCT GGG CGG CCC CGG CCC CGC 672
 Arg Glu Ile Leu Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg
 50 55 60

10

15

20

25

30

35

40

45

50

55

GCC AGT CCG AGT CCC ATC CGC ACC CCT CGG GCA GTG AGG CCA CTG AGG 1296
Ala Ser Pro Ser Pro Ile Arg Thr Pro Arg Ala Val Arg Pro Leu Arg
255 260 265

5 AGG AGG CAG CCG AAG AAA AGC AAC GAG CTG CCG CAG GCC AAC CGA CTC 1344
 Arg Arg Gln Pro Lys Lys Ser Asn Glu Leu Pro Gln Ala Asn Arg Leu
 270 275 280 285

10 CCA GGG ATC TTT GAT GAC GTC CAC GGC TCC CAC GGC CGG CAG GTC TGC 1392
 Pro Gly Ile Phe Asp Asp Val His Gly Ser His Gly Arg Gln Val Cys
 290 295 300

15 CGT CGG CAC GAG CTC TAC GTC AGC TTC CAG GAC CTC GGC TGG CTG GAC 1440
 Arg Arg His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Leu Asp
 305 310 315

20 TGG GTC ATC GCT CCC CAA GGC TAC TCG GCC TAT TAC TGT GAG GGG GAG 1488
 Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu
 320 325 330

25 TGC TCC TTC CCA CTG GAC TCC TGC ATG AAT GCC ACC AAC CAC GCC ATC 1536
 Cys Ser Phe Pro Leu Asp Ser Cys Met Asn Ala Thr Asn His Ala Ile
 335 340 345

30 CTG CAG TCC CTG GTG CAC CTG ATG AAG CCA AAC GCA GTC CCC AAG GCG 1584
 Leu Gln Ser Leu Val His Leu Met Lys Pro Asn Ala Val Pro Lys Ala
 350 355 360 365

35 TGC TGT GCA CCC ACC AAG CTG AGC GCC ACC TCT GTG CTC TAC TAT GAC 1632
 Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp
 370 375 380

40 AGC AGC AAC AAC GTC ATC CTG CGC AAA CAC CGC AAC ATG GTG GTC AAG 1680
 Ser Ser Asn Asn Val Ile Leu Arg Lys His Arg Asn Met Val Val Lys
 385 390 395

45 GCC TGC GGC TGC CAC T GAGTCAGCCC GCCCAGCCCT ACTGCAG 1723
 Ala Cys Gly Cys His
 400

(2) INFORMATION FOR SEQ ID NO:21:

(i)SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 402 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii)MOLECULE TYPE: protein

(ix)FEATURE:

(A)OTHER INFORMATION: /product= "hOP2-PP"

(xi)SEQUENCE DESCRIPTION: SEQ ID NO:21:

Met Thr Ala Leu Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys
 1 5 10 15

Ala	Leu	Gly	Gly	Gly	Pro	Gly	Leu	Arg	Pro	Pro	Pro	Gly	Cys	Pro
			20				25						30	
Gln	Arg	Arg	Leu	Gly	Ala	Arg	Glu	Arg	Arg	Asp	Val	Gln	Arg	Glu
		35					40					45		Ile
Leu	Ala	Val	Leu	Gly	Leu	Pro	Gly	Arg	Pro	Arg	Pro	Arg	Ala	Pro
	50					55					60			Pro
Ala	Ala	Ser	Arg	Leu	Pro	Ala	Ser	Ala	Pro	Leu	Phe	Met	Leu	Asp
	65				70					75				80
Tyr	His	Ala	Met	Ala	Gly	Asp	Asp	Asp	Glu	Asp	Gly	Ala	Pro	Ala
				85					90					95
Arg	Arg	Leu	Gly	Arg	Ala	Asp	Leu	Val	Met	Ser	Phe	Val	Asn	Met
			100					105					110	Val
Glu	Arg	Asp	Arg	Ala	Leu	Gly	His	Gln	Glu	Pro	His	Trp	Lys	Glu
		115					120					125		Phe
Arg	Phe	Asp	Leu	Thr	Gln	Ile	Pro	Ala	Gly	Glu	Ala	Val	Thr	Ala
	130					135					140			Ala
Glu	Phe	Arg	Ile	Tyr	Lys	Val	Pro	Ser	Ile	His	Leu	Leu	Asn	Arg
	145				150					155				160
Leu	His	Val	Ser	Met	Phe	Gln	Val	Val	Gln	Glu	Gln	Ser	Asn	Arg
				165					170					175
Ser	Asp	Leu	Phe	Phe	Leu	Asp	Leu	Gln	Thr	Leu	Arg	Ala	Gly	Asp
			180					185					190	Glu
Gly	Trp	Leu	Val	Leu	Asp	Val	Thr	Ala	Ala	Ser	Asp	Cys	Trp	Leu
		195					200					205		Leu
Lys	Arg	His	Lys	Asp	Leu	Gly	Leu	Arg	Leu	Tyr	Val	Glu	Thr	Glu
	210					215					220			Asp
Gly	His	Ser	Val	Asp	Pro	Gly	Leu	Ala	Gly	Leu	Leu	Gly	Gln	Arg
	225				230					235				Ala
Pro	Arg	Ser	Gln	Gln	Pro	Phe	Val	Val	Thr	Phe	Phe	Arg	Ala	Ser
				245					250					255
Ser	Pro	Ile	Arg	Thr	Pro	Arg	Ala	Val	Arg	Pro	Leu	Arg	Arg	Gln
			260					265					270	
Pro	Lys	Lys	Ser	Asn	Glu	Leu	Pro	Gln	Ala	Asn	Arg	Leu	Pro	Gly
		275					280					285		Ile
Phe	Asp	Asp	Val	His	Gly	Ser	His	Gly	Arg	Gln	Val	Cys	Arg	Arg
	290						295				300			His

Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Leu Asp Trp Val Ile
305 310 315 320

Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu Cys Ser Phe
325 330 335

Pro Leu Asp Ser Cys Met Asn Ala Thr Asn His Ala Ile Leu Gln Ser
340 345 350

Leu Val His Leu Met Lys Pro Asn Ala Val Pro Lys Ala Cys Cys Ala
355 360 365

Pro Thr Lys Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp Ser Ser Asn
370 375 380

Asn Val Ile Leu Arg Lys His Arg Asn Met Val Val Lys Ala Cys Gly
385 390 395 400

Cys His

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1926 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HURIDAE
- (F) TISSUE TYPE: EMBRYO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 93..1289
- (D) OTHER INFORMATION: /note= "mOP2 cDNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

GCCAGGCACA GGTGCGCCGT CTGGTCCTCC CCGTCTGGCG TCAGCCGAGC      50
CCGACCAGCT ACCAGTGGAT GCGCGCCGGC TGAAAGTCCG AG ATG GCT ATG CGT      104
                                     Met Ala Met Arg
                                     1
CCC GGG CCA CTC TGG CTA TTG GGC CTT GCT CTG TGC GCG CTG GGA GGC      152
Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys Ala Leu Gly Gly
5      5      10      15      20

```

5	GGC CAC GGT CCG CGT CCC CCG CAC ACC TGT CCC CAG CGT CGC CTG GGA Gly His Gly Pro Arg Pro Pro His Thr Cys Pro Gln Arg Arg Leu Gly 25 30 35	200
	GCG CGC GAG CGC CGC GAC ATG CAG CGT GAA ATC CTG GCG GTG CTC GGG Ala Arg Glu Arg Arg Asp Met Gln Arg Glu Ile Leu Ala Val Leu Gly 40 45 50	248
10	CTA CCG GGA CGG CCC CGA CCC CGT GCA CAA CCC GCG GCT GCC CGG CAG Leu Pro Gly Arg Pro Arg Pro Arg Ala Gln Pro Ala Ala Arg Gln 55 60 65	296
15	CCA GCG TCC GCG CCC CTC TTC ATG TTG GAC CTA TAC CAC GCC ATG ACC Pro Ala Ser Ala Pro Leu Phe Met Leu Asp Leu Tyr His Ala Met Thr 70 75 80	344
20	GAT GAC GAC GAC GGC GGG CCA CCA CAG GCT CAC TTA GGC CGT GCC GAC Asp Asp Asp Asp Gly Gly Pro Pro Gln Ala His Leu Gly Arg Ala Asp 85 90 95 100	392
	CTG GTC ATG AGC TTC GTC AAC ATG GTG GAA CGC GAC CGT ACC CTG GGC Leu Val Met Ser Phe Val Asn Met Val Glu Arg Asp Arg Thr Leu Gly 105 110 115	440
25	TAC CAG GAG CCA CAC TGG AAG GAA TTC CAC TTT GAC CTA ACC CAG ATC Tyr Gln Glu Pro His Trp Lys Glu Phe His Phe Asp Leu Thr Gln Ile 120 125 130	488
30	CCT GCT GGG GAG GCT GTC ACA GCT GCT GAG TTC CGG ATC TAC AAA GAA Pro Ala Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile Tyr Lys Glu 135 140 145	536
35	CCC AGC ACC CAC CCG CTC AAC ACA ACC CTC CAC ATC AGC ATG TTC GAA Pro Ser Thr His Pro Leu Asn Thr Thr Leu His Ile Ser Met Phe Glu 150 155 160	584
	GTG GTC CAA GAG CAC TCC AAC AGG GAG TCT GAC TTG TTC TTT TTG GAT Val Val Gln Glu His Ser Asn Arg Glu Ser Asp Leu Phe Phe Leu Asp 165 170 175 180	632
40	CTT CAG ACG CTC CGA TCT GGG GAC GAG GGC TGG CTG GTG CTG GAC ATC Leu Gln Thr Leu Arg Ser Gly Asp Glu Gly Trp Leu Val Leu Asp Ile 185 190 195	680
45	ACA GCA GCC AGT GAC CGA TGG CTG CTG AAC CAT CAC AAG GAC CTG GGA Thr Ala Ala Ser Asp Arg Trp Leu Leu Asn His His Lys Asp Leu Gly 200 205 210	728
50	CTC CGC CTC TAT GTG GAA ACC GCG GAT GGG CAC AGC ATC GAT CCT GGC Leu Arg Leu Tyr Val Glu Thr Ala Asp Gly His Ser Met Asp Pro Gly 215 220 225	776

	CTG GCT GGT CTG CTT GGA CGA CAA GCA CCA CGC TCC AGA CAG CCT TTC Leu Ala Gly Leu Leu Gly Arg Gln Ala Pro Arg Ser Arg Gln Pro Phe 230 235 240	824
5	ATG GTA ACC TTC TTC AGG GCC AGC CAG AGT CCT GTG CGG GCC CCT CGG Met Val Thr Phe Phe Arg Ala Ser Gln Ser Pro Val Arg Ala Pro Arg 245 250 255 260	872
10	GCA GCG AGA CCA CTG AAG AGG AGG CAG CCA AAG AAA ACG AAC GAG CTT Ala Ala Arg Pro Leu Lys Arg Arg Gln Pro Lys Lys Thr Asn Glu Leu 265 270 275	920
15	CCG CAC CCC AAC AAA CTC CCA GGG ATC TTT GAT GAT GGC CAC GGT TCC Pro His Pro Asn Lys Leu Pro Gly Ile Phe Asp Asp Gly His Gly Ser 280 285 290	968
	CGC GGC AGA GAG GTT TGC CGC AGG CAT GAG CTC TAC GTC AGC TTC CGT Arg Gly Arg Glu Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Arg 295 300 305	1016
20	GAC CTT GGC TGG CTG GAC TGG GTC ATC GCC CCC CAG GGC TAC TCT GCC Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala 310 315 320	1064
25	TAT TAC TGT GAG GGG GAG TGT GCT TTC CCA CTG GAC TCC TGT ATG AAC Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asp Ser Cys Met Asn 325 330 335 340	1112
	GCC ACC AAC CAT GCC ATC TTG CAG TCT CTG GTG CAC CTG ATG AAG CCA Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro 345 350 355	1160
30	GAT GTT GTC CCC AAG GCA TGC TGT GCA CCC ACC AAA CTG AGT GCC ACC Asp Val Val Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr 360 365 370	1208
35	TCT GTG CTG TAC TAT GAC AGC AGC AAC AAT GTC ATC CTG CGT AAA CAC Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His 375 380 385	1256
40	CGT AAC ATG GTG GTC AAG GCC TGT GGC TGC CAC TGAGGCCCG CCCAGCATCC Arg Asn Met Val Val Lys Ala Cys Gly Cys His 390 395	1309
	TGCTTCTACT ACCTTACCAT CTGGCCGGGC CCCTCTCCAG AGGCAGAAAC CCTTCTATGT	1369
45	TATCATAGCT CAGACAGGGG CAATGGGAGG CCCTTCACTT CCCCTGGCCA CTTCTGCTA	1429
	AAATTCTGGT CTTTCCCAGT TCCTCTGTCC TTCATGGGGT TTCGGGGCTA TCACCCCGCC	1489
	CTCTCCATCC TCCTACCCCA AGCATAGACT GAATGCACAC AGCATCCCAG AGCTATGCTA	1549

50

55

ACTGAGAGGT CTGGGGTCAG CACTGAAGGC CCACATGAGG AAGACTGATC CTTGGCCATC 1609
 CTCAGCCCAC AATGGCAAAT TCTGGATGGT CTAAGAAGGC CGTGGAATTC TAAACTAGAT 1669
 5 GATCTGGGCT CTCTGCACCA TTCATTGTGG CAGTTGGGAC ATTTTtagGT ATAACAGACA 1729
 CATACACTTA GATCAATGCA TCGCTGTACT CCTTGAAATC AGAGCTAGCT TGTTAGAAAA 1789
 AGAATCAGAG CCAGGTATAG CGGTGCATGT CATTAAATCCC AGCGCTAAAG AGACAGAGAC 1849
 10 AGGAGAATCT CTGTGAGTTC AAGGCCACAT AGAAAGAGCC TGTCTCGGGA GCAGGAAAAA 1909
 AAAAAAAAAAC GGAATTC 1926

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 399 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(D) OTHER INFORMATION: /product= "mOP2-PP"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met Ala Met Arg Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys
 1 5 10 15
 Ala Leu Gly Gly Gly His Gly Pro Arg Pro Pro His Thr Cys Pro Gln
 20 25 30
 Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Met Gln Arg Glu Ile Leu Ala
 35 40 45
 Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg Ala Gln Pro Ala Ala
 50 55 60 65
 Ala Arg Gln Pro Ala Ser Ala Pro Leu Phe Met Leu Asp Leu Tyr His Ala
 70 75 80
 Met Thr Asp Asp Asp Asp Gly Gly Pro Pro Gln Ala His Leu Gly Arg
 85 90 95
 Ala Asp Leu Val Met Ser Phe Val Asn Met Val Glu Arg Asp Arg Thr
 100 105 110
 Leu Gly Tyr Gln Glu Pro His Trp Lys Glu Phe His Phe Asp Leu Thr
 115 120 125 130

Gln Ile Pro Ala Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile Tyr
 135 140 145
 5 Lys Glu Pro Ser Thr His Pro Leu Asn Thr Thr Leu His Ile Ser Met
 150 155 160
 Phe Glu Val Val Gln Glu His Ser Asn Arg Glu Ser Asp Leu Phe Phe
 165 170 175
 10 Leu Asp Leu Gln Thr Leu Arg Ser Gly Asp Glu Gly Trp Leu Val Leu
 180 185 190
 Asp Ile Thr Ala Ala Ser Asp Arg Trp Leu Leu Asn His His Lys Asp
 195 200 205 210
 15 Leu Gly Leu Arg Leu Tyr Val Glu Thr Ala Asp Gly His Ser Met Asp
 215 220 225
 Pro Gly Leu Ala Gly Leu Leu Gly Arg Gln Ala Pro Arg Ser Arg Gln
 230 235 240
 Pro Phe Met Val Thr Phe Phe Arg Ala Ser Gln Ser Pro Val Arg Ala
 245 250 255
 25 Pro Arg Ala Ala Arg Pro Leu Lys Arg Arg Gln Pro Lys Lys Thr Asn
 260 265 270
 Glu Leu Pro His Pro Asn Lys Leu Pro Gly Ile Phe Asp Asp Gly His
 275 280 285 290
 30 Gly Ser Arg Gly Arg Glu Val Cys Arg Arg His Glu Leu Tyr Val Ser
 295 300 305
 Phe Arg Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr
 310 315 320
 35 Ser Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asp Ser Cys
 325 330 335
 Met Asn Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His Leu Met
 340 345 350
 Lys Pro Asp Val Val Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser
 355 360 365 370
 45 Ala Thr Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg
 375 380 385
 Lys His Arg Asn Met Val Val Lys Ala Cys Gly Cys His
 390 395
 50

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1368 base pairs
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1368

(D) OTHER INFORMATION:/STANDARD NAME="60A"

(x) PUBLICATION INFORMATION:

(A) AUTHORS: WHARTON, KRISTI A.; THOMSEN, GERALD H.; GELBERT, WILLIAM M.

(B) TITLE: DROSOPHILA 60A GENE...

(C) JOURNAL: PROC. NAT'L ACAD. SCI. USA

(D) VOLUME: 88

(E) RELEVANT RESIDUES IN SEQ ID NO:3: FROM 1 TO 1368

(F) PAGES: 9214-9218

(G) DATE: OCT - 1991

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

ATG TCG GGA CTG CGA AAC ACC TCG GAG GCC GTT GCA GTG CTC GCC TCC	48
Met Ser Gly Leu Arg Asn Thr Ser Glu Ala Val Ala Val Leu Ala Ser	
1 5 10 15	
CTG GGA CTC GGA ATG GTT CTG CTC ATG TTC GTG GCG ACC ACG CCG CCG	96
Leu Gly Leu Gly Met Val Leu Leu Met Phe Val Ala Thr Thr Pro Pro	
20 25 30	
GCC GTT GAG GCC ACC CAG TCG GGG ATT TAC ATA GAC AAC GGC AAG GAC	144
Ala Val Glu Ala Thr Gln Ser Gly Ile Tyr Ile Asp Asn Gly Lys Asp	
35 40 45	
CAG ACG ATC ATG CAC AGA GTG CTG AGC GAG GAC GAC AAG CTG GAC GTC	192
Gln Thr Ile Met His Arg Val Leu Ser Glu Asp Asp Lys Leu Asp Val	
50 55 60	
TCG TAC GAG ATC CTC GAG TTC CTG GGC ATC GCC GAA CGG CCG ACG CAC	240
Ser Tyr Glu Ile Leu Glu Phe Leu Gly Ile Ala Glu Arg Pro Thr His	
65 70 75 80	
CTG AGC AGC CAC CAG TTG TCG CTG AGG AAG TCG GCT CCC AAG TTC CTG	288
Leu Ser Ser His Gln Leu Ser Leu Arg Lys Ser Ala Pro Lys Phe Leu	
85 90 95	
CTG GAC GTC TAC CAC CGC ATC ACG GCG GAG GAG GGT CTC AGC GAT CAG	336
Leu Asp Val Tyr His Arg Ile Thr Ala Glu Glu Gly Leu Ser Asp Gln	
100 105 110	

5	GAT GAG GAC GAC GAC TAC GAA CGC GGC CAT CGG TCC AGG AGG AGC GCC Asp Glu Asp Asp Asp Tyr Glu Arg Gly His Arg Ser Arg Arg Ser Ala 115 120 125	384
10	GAC CTC GAG GAG GAT GAG GGC GAG CAG CAG AAG AAC TTC ATC ACC GAC Asp Leu Glu Glu Asp Glu Gly Glu Gln Gln Lys Asn Phe Ile Thr Asp 130 135 140	432
15	CTG GAC AAG CGG GCC ATC GAC GAG AGC GAC ATC ATC ATG ACC TTC CTG Leu Asp Lys Arg Ala Ile Asp Glu Ser Asp Ile Ile Met Thr Phe Leu 145 150 155 160	480
20	AAC AAG CGC CAC CAC AAT GTG GAC GAA CTG CGT CAC GAG CAC GGC CGT Asn Lys Arg His His Asn Val Asp Glu Leu Arg His Glu His Gly Arg 165 170 175	528
25	CGC CTG TGG TTC GAC GTC TCC AAC GTG CCC AAC GAC AAC TAC CTG GTG Arg Leu Trp Phe Asp Val Ser Asn Val Pro Asn Asp Asn Tyr Leu Val 180 185 190	576
30	ATG GCC GAG CTG CGC ATC TAT CAG AAC GCC AAC GAG GGC AAG TGG CTG Met Ala Glu Leu Arg Ile Tyr Gln Asn Ala Asn Glu Gly Lys Trp Leu 195 200 205	624
35	ACC GCC AAC AGG GAG TTC ACC ATC ACG GTA TAC GCC ATT GGC ACC GGC Thr Ala Asn Arg Glu Phe Thr Ile Thr Val Tyr Ala Ile Gly Thr Gly 210 215 220	672
40	ACG CTG GGC CAG CAC ACC ATG GAG CCG CTG TCC TCG GTG AAC ACC ACC Thr Leu Gly Gln His Thr Met Glu Pro Leu Ser Val Asn Thr Thr 225 230 235 240	720
45	GGG GAC TAC GTG GGC TGG TTG GAG CTC AAC GTG ACC GAG GGC CTG CAC Gly Asp Tyr Val Gly Trp Leu Glu Leu Asn Val Thr Glu Gly Leu His 245 250 255	768
50	GAG TGG CTG GTC AAG TCG AAG GAC AAT CAT GGC ATC TAC ATT GGA GCA Glu Trp Leu Val Lys Ser Lys Asp Asn His Gly Ile Tyr Ile Gly Ala 260 265 270	816
55	CAC GCT GTC AAC CGA CCC GAC CGC GAG GTG AAG CTG GAC GAC ATT GGA His Ala Val Asn Arg Pro Asp Arg Glu Val Lys Leu Asp Asp Ile Gly 275 280 285	864
60	CTG ATC CAC CGC AAG GTG GAC GAC GAG TTC CAG CCC TTC ATG ATC GGC Leu Ile His Arg Lys Val Asp Asp Glu Phe Gln Pro Phe Met Ile Gly 290 295 300	912
65	TTC TTC CGC GGA CCG GAG CTG ATC AAG GCG ACG GCC CAC AGC AGC CAC Phe Phe Arg Gly Pro Glu Leu Ile Lys Ala Thr Ala His Ser Ser His 305 310 315 320	960

CAC AGG AGC AAG CGA AGC GCC AGC CAT CCA CGC AAG CGC AAG AAG TCG 1008
 His Arg Ser Lys Arg Ser Ala Ser His Pro Arg Lys Arg Lys Lys Ser
 325 330 335

5 GTG TCG CCC AAC AAC GTG CCG CTG CTG GAA CCG ATG GAG AGC ACG CGC 1056
 Val Ser Pro Asn Asn Val Pro Leu Leu Glu Pro Met Glu Ser Thr Arg
 340 345 350

10 AGC TGC CAG ATG CAG ACC CTG TAC ATA GAC TTC AAG GAT CTG GGC TGG 1104
 Ser Cys Gln Met Gln Thr Leu Tyr Ile Asp Phe Lys Asp Leu Gly Trp
 355 360 365

15 CAT GAC TGG ATC ATC GCA CCA GAG GGC TAT GGC GCC TTC TAC TGC AGC 1152
 His Asp Trp Ile Ile Ala Pro Glu Gly Tyr Gly Ala Phe Tyr Cys Ser
 370 375 380

GGC GAG TGC AAT TTC CCG CTC AAT GCG CAC ATG AAC GCC ACG AAC CAT 1200
 Gly Glu Cys Asn Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His
 385 390 395 400

20 GCG ATC GTC CAG ACC CTG GTC CAC CTG CTG GAG CCC AAG AAG GTG CCC 1248
 Ala Ile Val Gln Thr Leu Val His Leu Leu Glu Pro Lys Lys Val Pro
 405 410 415

25 AAG CCC TGC TGC GCT CCG ACC AGG CTG GGA GCA CTA CCC GTT CTG TAC 1296
 Lys Pro Cys Cys Ala Pro Thr Arg Leu Gly Ala Leu Pro Val Leu Tyr
 420 425 430

30 CAC CTG AAC GAC GAG AAT GTG AAC CTG AAA AAG TAT AGA AAC ATG ATT 1344
 His Leu Asn Asp Glu Asn Val Asn Leu Lys Lys Tyr Arg Asn Met Ile
 435 440 445

GTG AAA TCC TGC GGG TGC CAT TGA 1368
 Val Lys Ser Cys Gly Cys His
 450 455

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 455 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met Ser Gly Leu Arg Asn Thr Ser Glu Ala Val Ala Val Leu Ala Ser
 1 5 10 15

Leu Gly Leu Gly Met Val Leu Leu Met Phe Val Ala Thr Thr Pro Pro
 20 25 30

Ala Val Glu Ala Thr Gln Ser Gly Ile Tyr Ile Asp Asn Gly Lys Asp
 35 40 45
 5 Gln Thr Ile Met His Arg Val Leu Ser Glu Asp Asp Lys Leu Asp Val
 50 55 60
 Ser Tyr Glu Ile Leu Glu Phe Leu Gly Ile Ala Glu Arg Pro Thr His
 65 70 75 80
 10 Leu Ser Ser His Gln Leu Ser Leu Arg Lys Ser Ala Pro Lys Phe Leu
 85 90 95
 Leu Asp Val Tyr His Arg Ile Thr Ala Glu Glu Gly Leu Ser Asp Gln
 100 105 110
 15 Asp Glu Asp Asp Asp Tyr Glu Arg Gly His Arg Ser Arg Arg Ser Ala
 115 120 125
 Asp Leu Glu Glu Asp Glu Gly Glu Gln Gln Lys Asn Phe Ile Thr Asp
 130 135 140
 20 Leu Asp Lys Arg Ala Ile Asp Glu Ser Asp Ile Ile Met Thr Phe Leu
 145 150 155 160
 25 Asn Lys Arg His His Asn Val Asp Glu Leu Arg His Glu His Gly Arg
 165 170 175
 Arg Leu Trp Phe Asp Val Ser Asn Val Pro Asn Asp Asn Tyr Leu Val
 180 185 190
 30 Met Ala Glu Leu Arg Ile Tyr Gln Asn Ala Asn Glu Gly Lys Trp Leu
 195 200 205
 Thr Ala Asn Arg Glu Phe Thr Ile Thr Val Tyr Ala Ile Gly Thr Gly
 210 215 220
 35 Thr Leu Gly Gln His Thr Met Glu Pro Leu Ser Ser Val Asn Thr Thr
 225 230 235 240
 40 Gly Asp Tyr Val Gly Trp Leu Glu Leu Asn Val Thr Glu Gly Leu His
 245 250 255
 Glu Trp Leu Val Lys Ser Lys Asp Asn His Gly Ile Tyr Ile Gly Ala
 260 265 270
 45 His Ala Val Asn Arg Pro Asp Arg Glu Val Lys Leu Asp Asp Ile Gly
 275 280 285
 Leu Ile His Arg Lys Val Asp Asp Glu Phe Gln Pro Phe Met Ile Gly
 290 295 300
 50 Phe Phe Arg Gly Pro Glu Leu Ile Lys Ala Thr Ala His Ser Ser His
 305 310 315 320
 55

His Arg Ser Lys Arg Ser Ala Ser His Pro Arg Lys Arg Lys Lys Ser
 325 330 335
 5 Val Ser Pro Asn Asn Val Pro Leu Leu Glu Pro Met Glu Ser Thr Arg
 340 345 350
 Ser Cys Gln Met Gln Thr Leu Tyr Ile Asp Phe Lys Asp Leu Gly Trp
 355 360 365
 10 His Asp Trp Ile Ile Ala Pro Glu Gly Tyr Gly Ala Phe Tyr Cys Ser
 370 375 380
 Gly Glu Cys Asn Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His
 385 390 395 400
 15 Ala Ile Val Gln Thr Leu Val His Leu Leu Glu Pro Lys Lys Val Pro
 405 410 415
 Lys Pro Cys Cys Ala Pro Thr Arg Leu Gly Ala Leu Pro Val Leu Tyr
 420 425 430
 20 His Leu Asn Asp Glu Asn Val Asn Leu Lys Lys Tyr Arg Asn Met Ile
 435 440 445
 25 Val Lys Ser Cys Gly Cys His
 450 455

(2) INFORMATION FOR SEQ ID NO:26:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: protein
 (iii) ORIGINAL SOURCE:
 40 (A) ORGANISM: Homo Sapiens
 (ix) FEATURE:
 (A) NAME/KEY: Protein
 45 (B) LOCATION: 1..102
 (D) OTHER INFORMATION: /note="BHP3"

(2) INFORMATION FOR SEQ ID NO:26:

50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 55 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein

(ix)FEATURE:

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..104
- (D) OTHER INFORMATION: /note="BMP3"

(xi)SEQUENCE DESCRIPTION: SEQ ID NO:26:

Cys Ala Arg Arg Tyr Leu Lys Val Asp Phe Ala Asp Ile Gly Trp Ser
 1 5 10 15
 Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp Ala Tyr Try Cys Ser Gly
 20 25 30
 Ala Cys Gln Phe Pro Met Pro Lys Ser Leu Lys Pro Ser Asn His Ala
 35 40 45
 Thr Ile Gln Ser Ile Val Ala Arg Ala Val Gly Val Val Pro Gly Ile
 50 55 60
 Pro Glu Pro Cys Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu
 65 70 75 80
 Phe Phe Asp Glu Asn Lys Asn Val Val Leu Lys Val Tyr Pro Asn Met
 85 90 95
 Thr Val Glu Ser Cys Ala Cys Arg
 100

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 102 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HOMO SAPIENS

(ix) FEATURE:

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..102
- (D) OTHER INFORMATION: /note= "BHP5"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg Asp Leu Gly Trp Gln
 1 5 10 15

Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala Phe Tyr Cys Asp Gly
 20 25 30
 5 Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
 35 40 45
 10 Ile Val Gln Thr Leu Val His Leu Met Phe Pro Asp His Val Pro Lys
 50 55 60
 15 Pro Cys Cys Ala Pro Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe
 65 70 75 80
 Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val
 85 90 95
 Arg Ser Cys Gly Cys His
 100

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 102 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HOMO SAPIENS

(ix) FEATURE:

- (A) NAME/KEY: Protein
 (B) LOCATION: 1..102
 (D) OTHER INFORMATION: /note= "BMP6"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Cys Arg Lys His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Gln
 1 5 10 15
 Asp Trp Ile Ile Ala Pro Lys Gly Tyr Ala Ala Asn Tyr Cys Asp Gly
 20 25 30
 50 Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
 35 40 45
 Ile Val Gln Thr Leu Val His Leu Met Asn Pro Glu Tyr Val Pro Lys
 50 55 60
 55 Pro Cys Cys Ala Pro Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe
 65 70 75 80

Asp Asp Asn Ser Asn Val Ile Leu Lys Lys Tyr Arg Trp Met Val Val
 85 90 95

Arg Ala Cys Gly Cys His
 100

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 102 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: Protein
 (B) LOCATION: 1..102
 (D) OTHER INFORMATION: /label= OPX /note= "WHEREIN XAA AT EACH POS'N IS INDEPENDENTLY
 SELECTED FROM THE RESIDUES OCCURRING AT THE CORRESPONDING POS'N IN THE C-TER-
 HINAL SEQUENCE OF MOUSE OR HUMAN OP1 OR OP2 (SEE SEQ. ID NOS. 5,6,7 and 8 or 16,18,20
 and 22.)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Cys Xaa Xaa His Glu Leu Tyr Val Xaa Phe Xaa Asp Leu Gly Trp Xaa
 1 5 10 15

Asp Trp Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly
 20 25 30

Glu Cys Xaa Phe Pro Leu Xaa Ser Xaa Met Asn Ala Thr Asn His Ala
 35 40 45

Ile Xaa Gln Xaa Leu Val His Xaa Xaa Xaa Pro Xaa Xaa Val Pro Lys
 50 55 60

Xaa Cys Cys Ala Pro Thr Xaa Leu Xaa Ala Xaa Ser Val Leu Tyr Xaa
 65 70 75 80

Asp Xaa Ser Xaa Asn Val Xaa Leu Xaa Lys Xaa Arg Asn Met Val Val
 85 90 95

Xaa Ala Cys Gly Cys His
 100

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
 (B) TYPE: amino acids
 (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- 5 (A) NAME: Generic Sequence 5
 (D) OTHER INFORMATION: wherein each Xaa is independently selected from a group of one or more specified amino acids as defined in the specification.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

10

Leu Xaa Xaa Xaa Phe

1 5

15

Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa

10

Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala

15 20

20

Xaa Tyr Cys Xaa Gly Xaa Cys Xaa

25 30

Xaa Pro Xaa Xaa Xaa Xaa Xaa

35

25

Xaa Xaa Xaa Asn His Ala Xaa Xaa

40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa

50

30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys

55 60

35

Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa

65

40

Xaa Xaa Xaa Leu Xaa Xaa Xaa

70 75

45

Xaa Xaa Xaa Xaa Val Xaa Leu Xaa

80

Xaa Xaa Xaa Xaa Met Xaa Val Xaa

85 90

50

Xaa Cys Xaa Cys Xaa

95

(2) INFORMATION FOR SEQ ID NO:31:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102 amino acids

(B) TYPE: amino acids

(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME: Generic Sequence 6

(D) OTHER INFORMATION: wherein each Xaa is independently selected from a group of one or more specified amino acids as defined in the specification.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

```

Cys Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Phe
 1             5             10
Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
          15
Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala
20             25

Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
          30             35
Xaa Pro Xaa Xaa Xaa Xaa Xaa
          40
Xaa Xaa Xaa Asn His Ala Xaa Xaa
          45             50
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
          55
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
40             60             65
Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
          70
Xaa Xaa Xaa Leu Xaa Xaa Xaa
45             75             80
Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
          85
Xaa Xaa Xaa Xaa Met Xaa Val Xaa
50             90             95
Xaa Cys Xaa Cys Xaa
          100

```

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1238 base pairs, 372 amino acids
- (B) TYPE: nucleic acid, amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) ORIGINAL SOURCE:

- (A) ORGANISM: human
- (F) TISSUE TYPE: BRAIN

(iv) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION:
- (D) OTHER INFORMATION:

/product= "GDF-1"
/note= "GDF-1 CDNA"

(x) PUBLICATION INFORMATION:

- (A) AUTHORS: Lee, Se-Jin
- (B) TITTLE: Expression of Growth/Differentiation Factor 1
- (C) JOURNAL: Proc. Nat'l Acad. Sci.
- (D) VOLUME: 88
- (E) RELEVANT RESIDUES: 1-1238
- (F) PAGES: 4250-4254
- (G) DATE: May-1991

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

	GGGGACACCG	CCCCGCCCT	CAGCCCCTG	GTCCCGGGCC	GCCGCGGACC	CTGCGCACTC	60
	TCTGGTCATC	GCCTGGGAGG	AAG ATG CCA CCG CCG CAG CAA GGT CCC TGC GGC	113			
5			Met Pro Pro Pro Gln Gln Gly Pro Cys Gly				
		1	5	10			
	CAC CAC CTC CTC CTC CTC CTG GCC CTG CTG CTG CCC TCG CTG CCC	158					
	His His Leu Leu Leu Leu Leu Ala Leu Leu Leu Pro Ser Leu Pro						
		15	20	25			
10	CTG ACC CGC GCC CCC GTG CCC CCA GGC CCA GCC GCC GCC CTG CTC	203					
	Leu Thr Arg Ala Pro Val Pro Pro Gly Pro Ala Ala Ala Leu Leu						
		30	35	40			
15	CAG GCT CTA GGA CTG CGC GAT GAG CCC CAG GGT GCC CCC AGG CTC	248					
	Gln Ala Leu Gly Leu Arg Asp Glu Pro Gln Gly Ala Pro Arg Leu						
		45	50	55			
	CGG CCG GTT CCC CCG GTC ATG TGG CGC CTG TTT CGA CGC CGG GAC	293					
	Arg Pro Val Pro Pro Val Met Trp Arg Leu Phe Arg Arg Arg Asp						
20		60	65	70			
	CCC CAG GAG ACC AGG TCT GGC TCG CGG CGG ACG TCC CCA GGG GTC	338					
	Pro Gln Glu Thr Arg Ser Gly Ser Arg Arg Thr Ser Pro Gly Val						
		75	80	85			
25	ACC CTG CAA CCG TGC CAC GTG GAG GAG CTG GGG GTC GCC GGA AAC	383					
	Thr Leu Gln Pro Cys His Val Glu Glu Leu Gly Val Ala Gly Asn						
		90	95	100			
30	ATC GTG CGC CAC ATC CCG GAC CGC GGT GCG CCC ACC CGG GCC TCG	428					
	Ile Val Arg His Ile Pro Asp Arg Gly Ala Pro Thr Arg Ala Ser						
		105	110	115			
	GAG CCT GTC TCG GCC GCG GGG CAT TGC CCT GAG TGG ACA GTC GTC	473					
	Glu Pro Val Ser Ala Ala Gly His Cys Pro Glu Trp Thr Val Val						
35		120	125	130			
	TTC GAC CTG TCG GCT GTG GAA CCC GCT GAG CGC CCG AGC CGG GCC	518					
	Phe Asp Leu Ser Ala Val Glu Pro Ala Glu Arg Pro Ser Arg Ala						
		135	140	145			
40							
45							
50							
55							

	CGC	CTG	GAG	CTG	CGT	TTC	GCG	GCG	GCG	GCG	GCA	GCC	CCG	GAG	563	
	Arg	Leu	Glu	Leu	Arg	Phe	Ala	Ala	Ala	Ala	Ala	Ala	Pro	Glu		
					150									160		
5																
	GGC	GGC	TGG	GAG	CTG	AGC	GTG	GCG	CAA	GCG	GGC	CAG	GGC	GCG	GGC	608
	Gly	Gly	Trp	Glu	Leu	Ser	Val	Ala	Gln	Ala	Gly	Gln	Gly	Ala	Gly	
					165										175	
10																
	GCG	GAC	CCC	GGG	CCG	GTG	CTG	CTC	CGC	CAG	TTG	GTG	CCC	GCC	CTG	653
	Ala	Asp	Pro	Gly	Pro	Val	Leu	Leu	Arg	Gln	Leu	Val	Pro	Ala	Leu	
					180										190	
15																
	GGG	CCG	CCA	GTG	CGC	GCG	GAG	CTG	CTG	GGC	GCC	GCT	TGG	GCT	CGC	698
	Gly	Pro	Pro	Val	Arg	Ala	Glu	Leu	Leu	Gly	Ala	Ala	Trp	Ala	Arg	
					195										205	
20																
	AAC	GCC	TCA	TGG	CCG	CGC	AGC	CTC	CGC	CTG	GCG	CTG	GCG	CTA	CGC	743
	Asn	Ala	Ser	Trp	Pro	Arg	Ser	Leu	Arg	Leu	Ala	Leu	Ala	Leu	Arg	
					210										220	
25																
	CCC	CGG	GCC	CCT	GCC	GCC	TGC	GCG	CGC	CTG	GCC	GAG	GCC	TCG	CTG	788
	Pro	Arg	Ala	Pro	Ala	Ala	Cys	Ala	Arg	Leu	Ala	Glu	Ala	Ser	Leu	
					225										235	
30																
	CTG	CTG	GTG	ACC	CTC	GAC	CCG	CGC	CTG	TGC	CAC	CCC	CTG	GCC	CGG	833
	Leu	Leu	Val	Thr	Leu	Asp	Pro	Arg	Leu	Cys	His	Pro	Leu	Ala	Arg	
					240										250	
35																
	CCG	CGG	CGC	GAC	GCC	GAA	CCC	GTG	TTG	GGC	GGC	GGC	CCC	GGG	GGC	878
	Pro	Arg	Arg	Asp	Ala	Glu	Pro	Val	Leu	Gly	Gly	Gly	Pro	Gly	Gly	
					255										265	
40																
	GCT	TGT	CGC	GCG	CGG	CGG	CTG	TAC	GTG	AGC	TTC	CGC	CAG	GTG	GGC	923
	Ala	Cys	Arg	Ala	Arg	Arg	Leu	Tyr	Val	Ser	Phe	Arg	Glu	Val	Gly	
					270										280	
45																
	TGG	CAC	CGC	TGG	GTC	ATC	GCG	CCG	CGC	CCC	TTC	CTG	GCC	AAC	TAC	968
	Trp	His	Arg	Trp	Val	Ile	Arg	Pro	Arg	Gly	Phe	Leu	Ala	Asn	Tyr	
					285										295	
50																
	TGC	CAG	GGT	CAG	TGC	GCG	CTG	CCC	GTC	GCG	CTG	TCG	GGG	TCC	GGG	1013
	Cys	Gln	Gly	Gln	Cys	Ala	Leu	Pro	Val	Ala	Leu	Ser	Gly	Ser	Gly	
					300										310	
55																
	GGG	CCG	CCG	GCG	CTC	AAC	CAC	GCT	GTG	CTG	CGC	GCG	CTC	ATG	CAC	1058
	Gly	Pro	Pro	Ala	Leu	Asn	His	Ala	Val	Leu	Arg	Ala	Leu	Met	His	
					315										325	
60																
	GCG	GCC	GCC	CCG	GGA	GCC	GCC	GAC	CTG	CCC	TGC	TGC	GTG	CCC	GCG	1103
	Ala	Ala	Ala	Pro	Gly	Ala	Ala	Asp	Leu	Pro	Cys	Cys	Val	Pro	Ala	
					330										340	

10

GTG	GTG	CTG	CGG	CAG	TAT	GAG	GAC	ATG	GTG	GTG	GAC	GAG	TGC	GGC	1193
Val	Val	Leu	Arg	Gln	Tyr	Glu	Asp	Met	Val	Val	Asp	Glu	Cys	Gly	
				360					365					370	

TGC CGC TAACCCGGGG CGGGCAGGGA CCCGGGCCCA ACAATAAATG CCGCGTGG 1238
Cys Arg
372

(34) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: human
(F) TISSUE TYPE: BRAIN

(ix) **FEATURE:**

- (A) NAME/KEY: CDS
(B) LOCATION:
(D) OTHER INFORMATION: /function= /product= "GDF-1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

[illegible]

	Arg Pro Val Pro	Pro Val Met Trp Arg	Leu Phe Arg Arg Arg Asp
	60	65	70
5	Pro Gln Glu Thr	Arg Ser Gly Ser Arg	Arg Thr Ser Pro Gly Val
	75	80	85
	Thr Leu Gln Pro	Cyc His Val Glu Glu	Leu Gly Val Ala Gly Asn
	90	95	100
10	Ile Val Arg His	Ile Pro Asp Arg Gly	Ala Pro Thr Arg Ala Ser
	105	110	115
	Glu Pro Val Ser	Ala Ala Gly His Cys	Pro Glu Trp Thr Val Val
	120	125	130
15	Phe Asp Leu Ser	Ala Val Glu Pro Ala	Glu Arg Pro Ser Arg Ala
	135	140	145
	Arg Leu Glu Leu	Arg Phe Ala Ala Ala	Ala Ala Ala Pro Glu
	150	155	160
20	Gly Gly Trp Glu	Leu Ser Val Ala Gln	Ala Gly Gln Gly Ala Gly
	165	170	175
	Ala Asp Pro Gly	Pro Val Leu Leu Arg	Gln Leu Val Pro Ala Leu
	180	185	190
	Gly Pro Pro Val	Arg Ala Glu Leu Leu	Gly Ala Ala Trp Ala Arg
	195	200	205
30	Asn Ala Ser Trp	Pro Arg Ser Leu Arg	Leu Ala Leu Ala Leu Arg
	210	215	220
	Pro Arg Ala Pro	Ala Ala Cys Ala Arg	Leu Ala Glu Ala Ser Leu
	225	230	235
35	Leu Leu Val Thr	Leu Asp Pro Arg Leu	Cys His Pro Leu Ala Arg
	240	245	250
	Pro Arg Arg Asp	Ala Glu Pro Val Leu	Gly Gly Gly Pro Gly Gly
	255	260	265
40	Ala Cys Arg Ala	Arg Arg Leu Tyr Val	Ser Phe Arg Glu Val Gly
	270	275	280
	Trp His Arg Trp	Val Ile Arg Pro Arg	Gly Phe Leu Ala Asn Tyr
	285	290	295
45	Cys Gln Gly Gln	Cys Ala Leu Pro Val	Ala Leu Ser Gly Ser Gly
	300	305	310
50	Gly Pro Pro Ala	Leu Asn His Ala Val	Leu Arg Ala Leu Met His
	315	320	325

1. Use of a morphogen in the manufacture of a medicament for the treatment of metabolic bone disease by oral administration, wherein said morphogen:

- wherein the medicament does not comprise vitamin D or another bone resorption inhibitor.

- wherein the medicament does not comprise vitamin D or another bone resorption inhibitor.

3. The use of claim 1 or 2 wherein said morphogen is in a soluble form.
4. The use of claim 3 wherein said soluble form comprises a morphogen associated with a morphogen pro-domain.
5. The use of any one of the preceding claims wherein said metabolic bone disorder is any of osteoporosis, osteomalacia and renal osteodystrophy.
6. The use of any one of the preceding claims wherein said metabolic bone disorder is caused by a nutritional or hormonal deficiency.
7. The use of claim 2, wherein the morphogen is formulated for injection.
8. The use of claim 7, wherein the morphogen is formulated for intravenous injection.

Patentansprüche

1. Verwendung eines Morphogens bei der Herstellung eines Medikaments zur Behandlung einer metabolischen Knochenkrankung durch orale Verabreichung, wobei das Morphogen:

- (i) endochondrale Knochenbildung in einem in vivo Knochenassay stimuliert; und
- (ii) eine Aminosäuresequenz umfasst, ausgewählt aus der Gruppe bestehend aus irgendeinem von: OPX Sequenz definiert durch SEQ ID NO:29, menschlichem OP-1 (SEQ ID NO: 5, 16 und 17), OP-1 von der Maus (SEQ ID NO: 6, 18 und 19), menschlichem OP-2 (SEQ ID NO: 7, 20 und 21), OP-2 von der Maus (SEQ ID NO: 8, 22 und 23), 60A (SEQ ID NO: 24), GDF-1 (SEQ ID NO: 14, 32 und 33), BMP2A (SEQ ID NO: 9), BMP2B (SEQ ID NO: 10), DPP (SEQ ID NO: 23), Vgl (SEQ ID NO: 12), Vgr-1 (SEQ ID NO: 13), BMP3 (SEQ ID NO: 26), BMP5 (SEQ ID NO: 27) und BMP6 (SEQ ID NO: 28); und
- (iii) zur Verabreichung in einer zum Erhöhen des Verhältnisses von Spongiosavolumen zu gesamten Knochenvolumen in dem Säuger effektiven Menge geeignet ist;

wobei das Medikament kein Vitamin D oder einen anderen Knochenresorptionsinhibitor enthält.

2. Verwendung eines Morphogens bei der Herstellung eines Medikaments zur Behandlung einer metabolischen Knochenkrankung durch parenterale Verabreichung, wobei das Morphogen:

- (i) endochondrale Knochenbildung in einem in vivo Knochenassay stimuliert; und
- (ii) eine Aminosäuresequenz umfasst, ausgewählt aus der Gruppe bestehend aus irgendeinem von: menschlichem OP-1 (SEQ ID NO: 5, 16 und 17), OP-1 von der Maus (SEQ ID NO: 6, 18 und 19), menschlichem OP-2 (SEQ ID NO: 7, 20 und 21), OP-2 von der Maus (SEQ ID NO: 8, 22 und 23), 60A (SEQ ID NO: 24), GDF-1 (SEQ ID NO: 14, 32 und 33), DPP (SEQ ID NO: 23), Vgl (SEQ ID NO: 12), BMP5 (SEQ ID NO: 27) und BMP6 (SEQ ID NO: 28); und
- (iv) zur Verabreichung in einer zum Erhöhen des Verhältnisses von Spongiosavolumen zu gesamten Knochenvolumen in dem Säuger effektiven Menge geeignet ist;

wobei das Medikament kein Vitamin D oder einen anderen Knochenresorptionsinhibitor enthält.

- 3. Verwendung nach Anspruch 1 oder 2, wobei das Morphogen in einer löslichen Form vorliegt.
- 4. Verwendung nach Anspruch 3, wobei die lösliche Form ein Morphogen umfasst, das mit einer morphogenen Pro-Domäne zusammenhängt.
- 5. Verwendung nach einem der vorstehenden Ansprüche, wobei die metabolische Knochenkrankheit eine von Osteoporose, Osteomalazie und Nieren-Osteodystrophie ist.
- 6. Verwendung nach einem der vorstehenden Ansprüche, wobei die metabolische Knochenkrankheit durch einen Ernährungs- oder Hormonmangel verursacht ist.
- 7. Verwendung nach Anspruch 2, wobei das Morphogen zur Injektion formuliert ist.
- 8. Verwendung nach Anspruch 7, wobei das Morphogen zur intravenösen Injektion formuliert ist.

Revendications

1. Utilisation d'un morphogène dans la préparation d'un médicament pour le traitement de la maladie du métabolisme osseux par administration orale, ledit morphogène :

- (i) stimulant la formation d'os endochondral dans un examen des os in vivo ; et
- (ii) comprenant une séquence d'acides aminés choisie parmi le groupe constitué par : une séquence OPX définie par SEQ ID NO : 29, la OP-1 humaine (SEQ ID NO : 5, 16 et 17), la OP-1 de souris (SEQ ID NO: 6, 18 et 19), la OP-2 humaine (SEQ ID NO: 7, 20 et 21), la OP-2 de souris (SEQ ID NO: 8, 22 et 23), 60A (SEQ ID NO: 24), GDF-1 (SEQ ID NO: 14, 32 et 33), BMP2A (SEQ ID NO: 9), BMP2B (SEQ ID NO: 10), DPP (SEQ ID NO: 23), Vgl (SEQ ID NO: 12), Vgr-1 (SEQ ID NO: 13), BMP3 (SEQ ID NO: 26), BMP5 (SEQ ID NO: 27),

et BMP6 (SEQ ID NO: 28) ; et

(iii) étant approprié pour une administration en une quantité efficace pour augmenter le volume d'os spongieux par rapport au volume osseux total dans ledit mammifère,

5 le médicament ne comprenant pas de vitamine D ou un autre inhibiteur de la résorption osseuse.

2. Utilisation d'un morphogène dans la préparation d'un médicament pour le traitement de la maladie du métabolisme osseux par administration parentérale, ledit morphogène :

10 (i) stimulant la formation d'os endochondral dans un examen des os in vivo ; et

(ii) comprenant une séquence d'acides aminés choisie parmi le groupe constitué par : une séquence OPX définie par SEQ ID NO : 29, la OP-1 humaine (SEQ ID NO : 5, 16 et 17), la OP-1 de souris (SEQ ID NO: 6, 18 et 19), la OP-2 humaine (SEQ ID NO: 7, 20 et 21), la OP-2 de souris (SEQ ID NO: 8, 22 et 23), 60A (SEQ ID NO: 24), GDF-1 (SEQ ID NO: 14, 32 et 33), BMP2A (SEQ ID NO: 9), BMP2B (SEQ ID NO: 10), DPP (SEQ ID NO: 23), Vgl (SEQ ID NO: 12), Vgr-1 (SEQ ID NO: 13), BMP3 (SEQ ID NO: 26), BMP5 (SEQ ID NO: 27), et BMP6 (SEQ ID NO: 28) ; et

15 (iii) étant approprié pour une administration en une quantité efficace pour augmenter le volume d'os spongieux par rapport au volume osseux total dans ledit mammifère,

20 le médicament ne comprenant pas de vitamine D ou un autre inhibiteur de la résorption osseuse.

3. Utilisation selon la revendication 1 ou 2, dans laquelle ledit morphogène est présent sous une forme soluble.

25 4. Utilisation selon la revendication 3, dans laquelle ladite forme soluble comprend un morphogène en association avec un prodomaine de morphogène.

5. Utilisation selon l'une quelconque des revendications précédentes, dans laquelle ledit trouble du métabolisme osseux représente un trouble quelconque choisi parmi le groupe comprenant l'ostéoporose, l'ostéomalacie et l'ostéodystrophie rénale.

30 6. Utilisation selon l'une quelconque des revendications précédentes, dans laquelle ledit trouble du métabolisme osseux est provoqué par une déficience nutritionnelle ou hormonale.

35 7. Utilisation selon la revendication 2, dans laquelle le morphogène est formulé pour injection.

8. Utilisation selon la revendication 2, dans laquelle le morphogène est formulé pour injection intraveineuse.

40

45

50

55




Fig. 1




Fig. 2

PROTEIN CONCENTRATION (ng/ml)		cAMP (PICOMOLE/WELL)	
		-PTH	+PTH
BACKGROUND		1.30	2.20
OP - 1	1.0	1.25	3.45
	10.0	1.30	3.80
	40.0	1.25	4.45
TGF - β	0.1	0.95	1.42
	1.0	0.83	1.25
	5.0	0.68	0.88

Fig. 3




Fig. 4




Fig. 5




Fig. 6A




Fig. 6B




Fig. 7




Fig. 9




Fig. 8A




Fig. 8B




Fig. 10A




Fig. 10B




Fig. 11




Fig. 12B




Fig. 12A